

EQUIVARIANT COMPACTIFICATIONS OF REDUCTIVE GROUPS

D. A. TIMASHEV

ABSTRACT. We study equivariant projective compactifications of reductive groups obtained by closing the image of a group in the space of operators of a projective representation. We describe the structure and the mutual position of their orbits under the action of the doubled group by left/right multiplications, the local structure in a neighborhood of a closed orbit, and obtain some conditions of normality and smoothness of a compactification. Our methods of research use the theory of equivariant embeddings of spherical homogeneous spaces and of reductive algebraic semigroups.

Bibliography: 36 items.

0. INTRODUCTION

Let G be a connected reductive complex algebraic group. We may regard G as a symmetric homogeneous space $(G \times G)/\text{diag } G$ under the group $G \times G$ acting by left/right multiplications. Equivariant embeddings and, in particular, compactifications (completions) of G were constructed and studied in a number of papers. A completion of $G = \text{PGL}_n(\mathbb{C})$ via “complete collineations” was constructed by Semple [Sem]. Later, Neretin generalized this construction to other classical groups (see survey [Ner]). Neretin’s compactification may be considered as a particular case of the “wonderful” completion of a symmetric space, and in particular, of a semisimple adjoint group [CP-I] (see §8). A more general class of regular completions was considered in [CP-II] with applications to the intersection theory on homogeneous spaces. The geometry and cohomology of equivariant compactifications were studied in [CP], [St1], [BCP], [LP], [St2], [Br4], [BP], [AB2]. The theory of equivariant embeddings of reductive groups is a particular case of the theory of spherical varieties [LV], [Kn], [Br3] (see §4). On the other hand, affine equivariant embeddings of reductive groups are nothing else, but reductive algebraic semigroups, which were intensively studied

Date: February 21, 2003.

1991 Mathematics Subject Classification. 14L30, 14M17, 52B20.

Key words and phrases. Reductive group, projective representation, compactification, algebraic semigroup, spherical variety, orbit, local structure, normality, smoothness.

Supported by CRDF grant RM1–2088 and by RFBR grants 01–01–00756, 02–01–06402.

in a series of papers by Putcha and Renner [Pu1], [Re1], [PR], [Pu2], [Re2], and also by Vinberg [Vin], Rittatore [Rit] and Alexeev–Brion [AB1].

The compactifications of Semple–Neretin and the “wonderful” completion of de Concini and Procesi [CP-I] were based on explicit constructions of embeddings in projective spaces. However, the general theory of equivariant embeddings [LV] followed another way: the description of embeddings and the study of their geometric properties goes on in terms of certain objects of discrete convex geometry (colored cones and fans) and of their combinatorics, see §4.

Here we come back to a constructive viewpoint and consider the following natural class of compactifications of G . Let $G : \mathbb{P}(V)$ be a faithful projective representation. Then G embeds in the space $\mathbb{P}(\text{End}(V))$ of projective linear operators, where $\text{End}(V)$ is the algebra of linear operators on V . The projective closure $X = \overline{G} \subseteq \mathbb{P}(\text{End}(V))$ is a $(G \times G)$ -equivariant projective compactification of G . In fact, X is completely determined by the set of highest weights of $G : V$.

Our goal is to extract from these data some geometric information about X including $(G \times G)$ -orbit structure (Theorem 8), local structure in a neighborhood of a closed orbit (Propositions 6,7), conditions of normality (§10) and smoothness (§11). The problem of normality is important, in particular, because we can apply a well-developed theory of spherical varieties to normal X . For instance, in this case all orbit closures (including X itself) have rational singularities [Po], [BI], there is an explicit description of the Picard group [Br1] and, in certain cases, of the cohomology ring [BCP], [LP], [St2], [Br4], there are vanishing theorems for higher cohomology of nef line bundles [BI], etc.

Now we explain the structure of the paper. Basic notation is fixed in §1. In §2 we prove a technical lemma on decomposing tensor powers of G -modules into irreducibles. In §3 we consider a particular case of Luna’s fundamental lemma, which is required in examining smoothness of X . A brief exposition of the theory of spherical varieties, which is applied to studying projective compactifications of G , is given in §4. As our compactification X is a priori a non-normal variety, it is important to study its normalization $\tilde{X} \rightarrow X$. Some properties of the normalization of a spherical variety are discussed in Proposition 1 and in §5 for the particular case of toric varieties.

Another important tool for the study of X is the local structure of the projective action described in [BLV]. It provides a way to construct transversal slices to closed orbits and to reduce the study of the geometry of an action to actions of Levi subgroups on affine subvarieties. These results are recalled in §6.

The group G as a spherical (even symmetric) homogeneous space under $G \times G$ is considered in §7. Here we compute all combinatorial data required for the theory of spherical embeddings. In §8 we proceed to the

study of X , first by describing the closed orbits and the local structure of X in their neighborhoods in terms of the weight polytope of $G : V$. We observe that transversal slices to closed orbits have the structure of algebraic semigroups, whose unit groups are Levi subgroups in G . These results allow to compute combinatorial data describing $\tilde{X}$, which in turn are applied to describing the orbit structure of X in §9.

Normality and smoothness of X is discussed in §§10,11. Due to the local nature of these properties, everything is reduced to the case, where X is replaced by a transversal slice to a closed orbit, which is a reductive algebraic semigroup. While an effective normality criterion for the general case requires information on decomposing tensor products of reductive group representations and on branching to Levi subgroups, one can formulate some necessary or sufficient conditions and in certain cases (e.g. for regular highest weights) even criteria of normality. A criterion of smoothness is given in Theorem 9.

In §12 we illustrate the results obtained above by the study of equivariant compactifications of simple algebraic groups in the spaces of projective linear operators of fundamental and adjoint representations. Some of the results obtained here are related to similar results of Putcha–Renner [PR], [Re2] (orbital decomposition), and of Faltings [Fal], Kannan [Kan] and de Concini [Con] (normality) for reductive algebraic semigroups.

The aim of this paper is twofold. Together with obtaining new results, we gather in this paper and generalize some known results on reductive group embeddings, which are scattered in the literature. Therefore we tried to make the exposition maximally self-contained by including proofs of some known assertions (see e.g. §7).

Remark. For simplicity, we work over the field $\mathbb{C}$ of complex numbers. However our approach is purely algebraic, and all results are valid over any algebraically closed base field of characteristic 0.

The author is grateful to A. Stoyanovsky, whose interest to the subject stimulated the preparation of this paper, and to G. Röhrle and M. Brion for some useful remarks, which helped to improve the original text.

1. NOTATION

G is a connected reductive complex algebraic group.

$B \supseteq T, U$ are a fixed Borel subgroup, a maximal torus, and a maximal unipotent subgroup in G .

$\Delta = \Delta_G$ is the root system of G relative to T .

$\Delta^+ \supseteq \Pi = \{\alpha_1, \dots, \alpha_l\}$ are the subsystems of positive and simple roots relative to B .

$\alpha^\vee$ is the coroot dual to α (i.e., a 1-parameter subgroup in T).

e_α is a root vector in $\mathfrak{g}$ corresponding to α .

$u_\alpha(t) = \exp(te_\alpha)$ is a root unipotent 1-parameter subgroup.

$W = W_G = N_G(T)/T$ is the Weyl group.

$s_\alpha \in W$ is the reflection corresponding to $\alpha \in \Delta$ (or its representative in $N_G(T)$).

$Q = \mathbb{Z}\Delta$ is the root lattice.

$\mathfrak{X} = \mathfrak{X}(T)$ is the character lattice of T .

$C = C_G \subseteq \mathfrak{X} \otimes \mathbb{Q}$ is the positive Weyl chamber. We use the same notation for the positive Weyl chamber in the dual space $\text{Hom}(\mathfrak{X}, \mathbb{Q})$ identified with $\mathfrak{X} \otimes \mathbb{Q}$ via a W -invariant inner product.

$\langle \cdot, \cdot \rangle$ is the pairing between elements of dual spaces.

$\mathfrak{X}^+ = \mathfrak{X}_G^+ = C \cap \mathfrak{X}$ is the semigroup of dominant weights.

$\omega_1, \dots, \omega_l$ are the fundamental weights.

$V(\lambda) = V_G(\lambda)$ is an irreducible representation of highest weight λ .

$v_\mu \in V(\lambda)$ is (any) eigenvector of T -weight μ . In particular, v_λ is a highest weight vector.

$M_{(\lambda)}$ is the isotypic component of highest weight λ in a G -module M .

$\mathcal{P} = \mathcal{P}(V)$ is the weight polytope of a linear representation $G : V$.

$\langle v \rangle \in \mathbb{P}(V)$ is the point in the projective space corresponding to a vector $v \in V$.

$\text{End}(V)$ is the algebra of linear operators on V .

$\mathbb{C}(X)$ is the field of rational functions on a variety X .

$\mathbb{C}[X]$ is the algebra of polynomial functions on an affine variety X .

$\tilde{X}$ is the normalization of X .

Lie algebras of algebraic groups are denoted by the respective lower-case Gothic letters. The signs “ $\times, \rtimes$ ” denote *almost* (semi)direct products of algebraic groups (we allow intersections in finite subgroups).

For any parabolic $P \subseteq G$, $P \supseteq T$, we denote by P^- the opposite parabolic (and we also write $P = P^+$), and by P_u the unipotent radical.

2. A RESULT FROM REPRESENTATION THEORY

Let $G : V$ be any rational linear representation.

Lemma 1 (cf. [AB2, Lemma 4.9]). *For any $\mu \in C \cap \mathcal{P}$ there is n such that the decomposition of $V^{\otimes n}$ contains a simple submodule $V(n\mu)$.*

Proof. Put $\mathcal{M} = \{\mu \in C \mid \exists n \in \mathbb{N} : V^{\otimes n} \hookrightarrow V(n\mu)\}$. Clearly, $\mathcal{M} \subseteq \mathcal{P}$. We have to prove $\mathcal{M} = C \cap \mathcal{P}$.

Observe that $\mathcal{M}$ is convex. Indeed, for any $\mu, \nu \in \mathcal{M}$ consider any convex combination $r\mu + s\nu$, $r = p/m$, $s = q/m$, $p, q \geq 0$, $p+q = m$. For some n we have $V^{\otimes n} \hookrightarrow V(n\mu)$, $V(n\nu)$, whence $V^{\otimes nm} \hookrightarrow V(n\mu)^{\otimes p} \otimes V(n\nu)^{\otimes q} \hookrightarrow V(nm(r\mu + s\nu))$. Therefore $r\mu + s\nu \in \mathcal{M}$.

Thus it suffices to prove $\mathcal{M} = C \cap \mathcal{P}$ for irreducible $V = V(\lambda)$, $\lambda \in \mathfrak{X}^+$. Furthermore, we may assume G to be semisimple.

First, $\lambda \in \mathcal{M}$. Secondly, $0 \in \mathcal{M}$, since $V^{\otimes n} \supset \bigwedge^n V = V(0)$ for $n = \dim V$.

Other vertices μ of $C \cap \mathcal{P}$ are the intersection points of the faces of $\mathcal{P}$ at λ , whose direction subspaces are spanned by simple root subsystems, with perpendicular faces of C . If $L \subset G$ is the respective Levi subgroup, then the face is of the form $\mathcal{P}_L = \mathcal{P} \cap (\lambda + \langle \Delta_L \rangle)$, and μ is the center of $\mathcal{P}_L$. Moreover, $V(\lambda)$ contains $V_L(\lambda)$, whose weight polytope is $\mathcal{P}_L$. Under restriction to the commutator subgroup of L , μ maps to 0. Hence $V(\lambda)^{\otimes n} \hookrightarrow V_L(n\mu)$. But an L -highest weight vector of weight $n\mu$ in $V(\lambda)^{\otimes n}$ is automatically a highest weight vector for G (adding $\alpha \in \Delta^+ \setminus \Delta_L^+$ moves the weight outside the weight polytope $n\mathcal{P}$), whence $\mu \in \mathcal{M}$. $\square$

3. LUNA'S FUNDAMENTAL LEMMA

The following result is an easy particular case of the fundamental lemma [Lu, II.2] in the étale slice theory [loc. cit.]. For convenience of a reader, we provide a proof.

Lemma 2. *Suppose $X \subseteq V$ is a smooth closed G -stable subvariety with a dense orbit, and $0 \in X$. Then the projection $\pi : X \rightarrow T_0X \subseteq V$ along a G -stable complementary subspace is an isomorphism.*

Proof. The map π is étale at 0. The set of points, where π is not étale, is closed and G -stable. As closed orbits are separated by polynomial invariants, $\{0\}$ is the unique closed orbit in X , hence π is étale. By Zariski's Main Theorem, π decomposes into an open immersion $X \hookrightarrow \overline{X}$ and a finite morphism $\overline{X} \rightarrow T_0X$, so that $\mathbb{C}[\overline{X}]$ is the integral closure of $\mathbb{C}[T_0X]$ in $\mathbb{C}[X]$. But $\overline{X}$ also has a dense orbit, whence a unique closed orbit, which means $\overline{X} \setminus X = \emptyset$. Therefore π is a finite étale covering. But $\pi^{-1}(0) = \{0\}$ (X has a unique G -fixed point), hence $\deg \pi = 1$, i.e., π is an isomorphism. $\square$

4. SPHERICAL VARIETIES

A normal G -variety X is called *spherical* if B has an open orbit on X . The more so, X has an open G -orbit, which can be identified with a homogeneous space G/H by choosing a base point. Thus X may be considered as an equivariant embedding of G/H . All equivariant embeddings of a spherical homogeneous space G/H are described in terms of combinatorics of certain objects from convex geometry (cones, polytopes) related to G/H . This theory is due to Luna and Vust [LV]. For a transparent exposition, see [Kn], [Br3].

The algebra of regular functions on a spherical homogeneous space G/H is multiplicity free, i.e., nonzero isotypic components $\mathbb{C}[G/H]_{(\lambda)} \cong V(\lambda)$ are simple G -modules.

Let $\mathfrak{X}(G/H)$ be the weight lattice of all rational B -eigenfunctions on G/H (which are determined by their weights uniquely up to proportionality). Consider the dual space $E = \text{Hom}(\mathfrak{X}(G/H), \mathbb{Q})$. Any (discrete $\mathbb{Q}$ -valued) valuation of $\mathbb{C}(G/H)$ determines by restriction to

the multiplicative subgroup of rational B -eigenfunctions a homomorphism $\mathfrak{X}(G/H) \rightarrow \mathbb{Q}$, i.e., a point in E .

Theorem 1 ([Kn, 2.8, 6.3], [Br3, 3.1, 4.2]). *The set of G -invariant valuations of $\mathbb{C}(G/H)$ maps to E injectively, and its image is a solid polyhedral cone $\mathcal{V}$ containing the image of the negative Weyl chamber under the natural projection to E .*

B -invariant divisors on G/H (i.e., irreducible components of the complement to the open B -orbit, there are finitely many of them) determine a finite set $\mathcal{D}$ of valuations of $\mathbb{C}(G/H)$ (the set of *colors*), but the map $\rho : \mathcal{D} \rightarrow E$ is no more injective in general.

Definition. A *colored cone* is a pair $(\mathcal{C}, \mathcal{F})$, where $\mathcal{F} \subseteq \mathcal{D}$, $\rho(\mathcal{F}) \not\subseteq 0$, and $\mathcal{C}$ is a strictly convex polyhedral cone generated by $\rho(\mathcal{F})$ and by finitely many vectors from $\mathcal{V}$, so that $\text{int } \mathcal{C}$ (the relative interior) intersects $\mathcal{V}$.

A *colored face* of the colored cone is a colored cone $(\mathcal{C}', \mathcal{F}')$, where $\mathcal{C}'$ is a face of $\mathcal{C}$ and $\mathcal{F}' = \rho^{-1}(\rho(\mathcal{F}) \cap \mathcal{C}')$.

A *colored fan* is a finite collection of colored cones $(\mathcal{C}_i, \mathcal{F}_i)$ closed under passing to a colored face and such that $\text{int } \mathcal{C}_i \cap \text{int } \mathcal{C}_j \cap \mathcal{V} = \emptyset$, $\forall i \neq j$ (i.e., the cones intersect in faces inside $\mathcal{V}$).

Theorem 2 ([Kn, 4.3], [Br3, 3.4]). *There is a bijection between all equivariant normal embeddings $X \hookrightarrow G/H$ and all colored fans. Furthermore, G -orbits $Y \subseteq X$ are in bijection with colored cones $(\mathcal{C}_Y, \mathcal{F}_Y)$ of a given colored fan. The set of colors $\mathcal{F}_Y$ corresponds to all B -stable divisors on G/H , whose closures contain Y , and G -stable divisors containing Y determine the generators of those edges of $\mathcal{C}_Y$ which do not intersect $\rho(\mathcal{F}_Y)$.*

The geometry of a spherical variety is determined by its fan.

Theorem 3 ([Kn, 4.2, 5.2], [Br3, 3.4]). *Suppose $X \hookrightarrow G/H$ is a spherical variety, and $Y, Y' \subseteq X$ are G -orbits. Then $\overline{Y} \subseteq \overline{Y'}$ iff $(\mathcal{C}_{Y'}, \mathcal{F}_{Y'})$ is a face of $(\mathcal{C}_Y, \mathcal{F}_Y)$. In particular, G/H corresponds to $(0, \emptyset)$, and Y is projective iff $\mathcal{C}_Y$ is a solid cone. X is complete iff its fan covers all $\mathcal{V}$.*

Every spherical variety is covered by *simple* open subvarieties, which contain a unique closed orbit. In particular, affine spherical varieties are simple. The colored fan of a simple variety is determined by the colored cone of the closed orbit and consists of all its colored faces.

Affine spherical varieties have a more explicit description.

Theorem 4 ([Kn, 7.7]). *G/H admits an affine embedding iff $\rho(\mathcal{D})$ is contained in an open half-space of E . There is a bijection between all equivariant normal affine embeddings $X \hookrightarrow G/H$ and all strictly convex polyhedral cones $\mathcal{C} \subset E$ generated by $\rho(\mathcal{D})$ and by finitely many vectors*

from $\mathcal{V}$. Further,

$$\mathbb{C}[X] = \bigoplus_{\lambda \in \mathfrak{X}(G/H) \cap \mathcal{C}^\vee} \mathbb{C}[G/H]_{(\lambda)}$$

and $\mathbb{C}[X]^U = \mathbb{C}[\mathfrak{X}(G/H) \cap \mathcal{C}^\vee]$ is the semigroup algebra of the semigroup of lattice points in the dual cone $\mathcal{C}^\vee$. The colored fan of X is the set of all colored cones of $(\mathcal{C}, \mathcal{D})$.

Remark. The pair $(\mathcal{C}, \mathcal{D})$ of Theorem 4 might not be a colored cone in the sense of the above definition: $\text{int } \mathcal{C}$ may have empty intersection with $\mathcal{V}$. However we may consider its colored faces in accordance with the definition. There exists a largest face $\mathcal{C}' \subseteq \mathcal{C}$ whose interior intersects $\mathcal{V}$. The respective colored face $(\mathcal{C}', \mathcal{F}')$ corresponds to the closed orbit and determines the colored fan of X .

Proposition 1. *Let $X \hookrightarrow G/H$ be a quasiprojective embedding of a spherical homogeneous space. Then the normalization map $\tilde{X} \rightarrow X$ is bijective on the set of G -orbits.*

Proof. If X is affine, then $\mathbb{C}[X] = \bigoplus_{\lambda \in S} \mathbb{C}[G/H]_{(\lambda)}$, where $S \subseteq \mathfrak{X}^+$ is a finitely generated semigroup, and $\mathbb{Z}S = \mathfrak{X}(G/H)$. The following objects are in bijective correspondence: G -orbits on X , G -stable closed irreducible subvarieties of X , G -stable prime ideals in $\mathbb{C}[X]$, the (some, but generally not all) respective T -stable prime ideals in $\mathbb{C}[X]^U = \mathbb{C}[S]$ (the semigroup algebra of S), (some) sets of weights of the form $S \cap (\Sigma \setminus \Sigma')$, where $\Sigma = \mathbb{Q}_+ S$, and Σ' is its face.

After normalization, S is replaced by $\mathfrak{X}(G/H) \cap \Sigma$, but the cone Σ , its faces etc. do not change.

The projective case is reduced to the affine case by passing to the affine cone over X and extending G by homotheties. The general case reduces to the projective case by taking the projective closure. $\square$

Example. Toric varieties are a particular case of spherical varieties. Here $G = B = T$ is a torus, and we may assume $H = \{e\}$, if we want, after replacing T by T/H . The lattice $\mathfrak{X}(T/H)$ coincides with the character lattice of T/H . Every T -invariant valuation of $\mathbb{C}(T/H)$ is proportional to a valuation given by the order in $t \rightarrow \infty$ of a function restricted to a 1-parameter subgroup $\gamma(t) \in T/H$. Its value at an eigenfunction of weight λ equals $\langle \gamma, \lambda \rangle$, whence $\mathcal{V} = E$. There are no colors, and colored cones become usual cones, and a fan is just a finite collection of strictly convex polyhedral cones intersecting along faces.

A simple toric variety X given by a cone $\mathcal{C}$ is affine, and its coordinate algebra $\mathbb{C}[X] = \mathbb{C}[\mathfrak{X}(T/H) \cap \mathcal{C}^\vee]$ is the semigroup algebra of the semigroup of lattice points in the dual cone.

A projective toric variety can be also defined by a polytope in $E^* = \mathfrak{X}(T/H) \otimes \mathbb{Q}$ which is dual to the fan, i.e., the fan consists of dual cones to the cones at all vertices of the polytope (and of all their faces).

5. THE POLYTOPE OF THE CLOSURE OF A TORUS ORBIT

Consider a linear representation $T : V$ and a vector $v \in V$. The *support* $\text{supp } v$ is the convex hull of all weights in the weight decomposition of v .

Proposition 2. *The polytope of the projective toric variety $\widetilde{\overline{T\langle v \rangle}}$ equals $-\text{supp } v$.*

Proof. Consider the weight decomposition $v = v_{\lambda_0} + \cdots + v_{\lambda_n}$, complete the set $v_{\lambda_0}, \dots, v_{\lambda_n}$ to a weight basis of V , and consider the dual coordinates x_λ (of weights $-\lambda$) as homogeneous coordinates in $\mathbb{P}(V)$. The variety $X = \widetilde{\overline{T\langle v \rangle}}$ is covered by affine charts $X_i = \{x_{\lambda_i} \neq 0\}$. To be definite, consider X_0 . Then $\mathbb{C}[X_0] = \mathbb{C}[x_{\lambda_1}/x_{\lambda_0}, \dots, x_{\lambda_n}/x_{\lambda_0}]$ is the semigroup algebra of the semigroup $S_0 = \mathbb{Z}_+(\lambda_0 - \lambda_1) + \cdots + \mathbb{Z}_+(\lambda_0 - \lambda_n)$. This semigroup generates the cone Σ_0 at the vertex $-\lambda_0$ of $-\text{supp } v$. Hence $\mathbb{C}[X_0] = \mathbb{C}[\mathbb{Q}S_0 \cap \Sigma_0]$. Thus the fan of $\widetilde{X}$ is dual to $-\text{supp } v$. $\square$

Corollary (of the proof). *$\widetilde{\overline{T\langle v \rangle}}$ is normal iff $\mathbb{Q}S_i \cap \Sigma_i = S_i$, $i = 0, \dots, n$*

6. THE LOCAL STRUCTURE OF A PROJECTIVE ACTION

Consider a linear representation $G : V$ and fix a lowest weight vector $v_\lambda \in V$. Let $P^- = P(\lambda) = G_{\langle v_\lambda \rangle}$, and $P \supseteq B$ be the opposite parabolic. Consider the Levi decompositions $P = P_u \ltimes L$, $P^- = P_u^- \ltimes L$. We have a decomposition $V = \mathfrak{p}_u v_\lambda \oplus \mathbb{C}v_\lambda \oplus M$ in a direct sum of L -modules. Consider the dual highest weight vector $v_{-\lambda} \in V^*$ and the open subset $\mathring{V} = \mathfrak{p}_u v_\lambda \oplus \mathbb{C}^\times v_\lambda \oplus M$, which is the complement to the hyperplane $\{\langle v_{-\lambda}, \cdot \rangle = 0\}$.

Theorem 5 ([BLV]). *The P -action on $\mathbb{C}v_\lambda \oplus M$ yields an isomorphism*

$$\mathbb{P}(\mathring{V}) \cong P \times_L M(-\lambda) = P_u \times M(-\lambda)$$

where $M(-\lambda) = M$ is equipped with the L -action twisted by the character $-\lambda$ and embedded in $\mathbb{P}(V)$ as $\{\langle v_\lambda \rangle\} \times M = \mathbb{P}(v_\lambda + M)$.

It follows that any projective G -variety $X \subseteq \mathbb{P}(V)$ containing the closed orbit $Y = G\langle v_\lambda \rangle$ contains the affine open P -stable subset $\mathring{X} = X \cap \mathbb{P}(\mathring{V}) \cong P \times_L Z = P_u \times Z$ intersecting Y , where $Z \subseteq M(-\lambda)$ is a closed affine L -subvariety.

If X is a spherical G -variety, then Z is an affine spherical L -variety. It is easily seen from the structure of $\mathring{X}$ that $\mathfrak{X}(X) = \mathfrak{X}(\mathring{X}) = \mathfrak{X}(Z)$, and $\mathcal{C}_Y = \mathcal{C}_{Y \cap Z}$. The only delicate point is that the cone of invariant valuations can increase and the set of colors can decrease, because some colors may become L -stable divisors. Combined with Theorem 4, this yields an effective description of the local structure of a spherical variety by its fan.

7. A REDUCTIVE GROUP AS A SPHERICAL HOMOGENEOUS SPACE

The group G is a homogeneous space under $G \times G$ acting by left and right multiplications with stabilizer $\text{diag } G$ of e . Fix a Borel subgroup $B^- \times B \subseteq G \times G$. The Bruhat decomposition implies that $G = (G \times G) / \text{diag } G$ is a spherical homogeneous space. The combinatorial data related to this space in the sense of §4 were computed by Vust [Vu] (in the more general context of symmetric spaces) and by Rittatore [Rit]. We reproduce their results below. The following result is well known [Kr, II.3.1, Satz 3]:

Proposition 3. $\mathbb{C}[G] = \bigoplus_{\lambda \in \mathfrak{X}^+} \mathbb{C}[G]_{(\lambda)}$, where $\mathbb{C}[G]_{(\lambda)} \cong V(\lambda)^* \otimes V(\lambda)$ is the linear span of the matrix elements of the representation $G : V(\lambda)$.

Corollary. $\mathfrak{X}(G) \cong \mathfrak{X}$

The eigenfunction $f_\lambda(g) = \langle v_{-\lambda}, gv_\lambda \rangle$ has highest weight $(-\lambda, \lambda)$, i.e., λ under our identifications.

Proposition 4. $\mathbb{C}[G]_{(\lambda)} \cdot \mathbb{C}[G]_{(\mu)} = \mathbb{C}[G]_{(\lambda+\mu)} \oplus \bigoplus_i \mathbb{C}[G]_{(\lambda+\mu-\beta_i)}$, where $\lambda + \mu - \beta_i$ are highest weights of all “lower” irreducible components in the decomposition $V(\lambda) \otimes V(\mu) = V(\lambda + \mu) \oplus \dots$, so that $\beta_i \in \mathbb{Z}_+ \Pi$.

Proof. $\mathbb{C}[G]_{(\lambda)} \cdot \mathbb{C}[G]_{(\mu)}$ is generated by products of matrix elements of $G : V(\lambda)$ and $G : V(\mu)$, i.e., by matrix elements of $G : V(\lambda) \otimes V(\mu)$. $\square$

Corollary. $\mathcal{V} = -C$ is the negative Weyl chamber.

Proof. It is easy to see that a G -invariant valuation of $\mathbb{C}(G)$ is constant at any isotypic component $\mathbb{C}[G]_{(\lambda)}$, and its restriction is a linear function of λ , i.e., $\nu \in E = \text{Hom}(\mathfrak{X}, \mathbb{Q})$. The value of the valuation at $\forall f \in \mathbb{C}[G]$ equals $\min_{f_{(\lambda)} \neq 0} \langle \nu, \lambda \rangle$, where $f_{(\lambda)}$ is the projection of f on $\mathbb{C}[G]_{(\lambda)}$. Multiplying functions from $\mathbb{C}[G]_{(\lambda)}$ and $\mathbb{C}[G]_{(\mu)}$, we deduce from Proposition 4 that $\langle \nu, \beta_i \rangle \leq 0$ is necessary for ν to define a valuation. Since β_i are positive combinations of simple roots, and multiples of all simple roots are among them for appropriate $\lambda, \mu \in \mathfrak{X}^+$ (see e.g. §2), these inequalities define $-C$.

Conversely, each $\nu \in -C$ defines a $\mathbb{Q}$ -valued function on $\mathbb{C}[G]$ (denoted by the same letter by abuse of notation) satisfying the additive property of a valuation by the above formula. To verify the multiplicative property, take $p, q \in \mathbb{C}[G]$ and choose $\gamma \in \text{int}(-C)$ such that $\min \langle \gamma, \lambda \rangle$ over all λ with $p_{(\lambda)} \neq 0$, $\langle \nu, \lambda \rangle = \min$, and $\min \langle \gamma, \mu \rangle$ over all μ with $q_{(\mu)} \neq 0$, $\langle \nu, \mu \rangle = \min$, are reached at the unique points λ_0 and μ_0 , respectively. Then $pq = f_{(\lambda_0+\mu_0)} + \sum f_{(\chi)}$, where $\langle \nu, \chi \rangle \geq \langle \nu, \lambda_0 + \mu_0 \rangle$, $\langle \gamma, \chi \rangle \geq \langle \gamma, \lambda_0 + \mu_0 \rangle$, and at least one of these inequalities is strict. Hence $\nu(pq) = \langle \nu, \lambda_0 + \mu_0 \rangle = \nu(p) + \nu(q)$, and we are done. $\square$

It follows from the Bruhat decomposition, that $(B^- \times B)$ -stable divisors on G are of the form $D_i = \overline{B^- s_{\alpha_i} B}$. If G is of simply connected type (i.e., G is a direct product of a torus and a simply connected

semisimple group), then $\omega_i \in \mathfrak{X}$, and D_i is defined by the equation $f_{\omega_i} = 0$. More precisely, consider a curve $g_j(t) = u_{-\alpha_j}(t)\alpha_j^\vee(t^{-1})u_{\alpha_j}(-t)$ in the big Bruhat cell. Then $\lim_{t \rightarrow \infty} g_j(t) = s_{\alpha_j} \in D_j$ (everything takes place in an SL_2 -subgroup, where it is computed explicitly), and

$$f_{\omega_i}(g_j(t)) = \langle v_{-\omega_i}, \alpha_j^\vee(t^{-1})v_{\omega_i} \rangle = t^{-\langle \omega_i, \alpha_j^\vee \rangle} \langle v_{-\omega_i}, v_{\omega_i} \rangle$$

Hence f_{ω_i} has order 1 along D_j for $i = j$ and 0 for $i \neq j$. Therefore the colors look like $\rho(D_j) = \alpha_j^\vee$. This conclusion remains valid for arbitrary G , because G is covered by a group of simply connected type with the colors being the preimages of the colors of G .

These results, due to Vust [Vu], allow us to apply the theory of §§4,6 to describing $(G \times G)$ -equivariant embeddings of G . Affine embeddings have another remarkable property.

Theorem 6 ([Vin]). *An affine $(G \times G)$ -equivariant embedding $X \hookrightarrow G$ is an algebraic semigroup with unit, and G is the group of invertibles in X .*

Proof. The actions of the left and right copy of $G \times G$ on X define coactions $\mathbb{C}[X] \rightarrow \mathbb{C}[G] \otimes \mathbb{C}[X]$ and $\mathbb{C}[X] \rightarrow \mathbb{C}[X] \otimes \mathbb{C}[G]$, which are the restrictions to $\mathbb{C}[X] \subseteq \mathbb{C}[G]$ of the comultiplication $\mathbb{C}[G] \rightarrow \mathbb{C}[G] \otimes \mathbb{C}[G]$. Hence the image of $\mathbb{C}[X]$ lies in $\mathbb{C}[G] \otimes \mathbb{C}[X] \cap \mathbb{C}[X] \otimes \mathbb{C}[G] = \mathbb{C}[X] \otimes \mathbb{C}[X]$, and we have a comultiplication in $\mathbb{C}[X]$. Now G is open in X and consists of invertibles. For any invertible $x \in X$, we have $xG \cap G \neq \emptyset \implies x \in G$. $\square$

8. PROJECTIVE COMPACTIFICATION OF A REDUCTIVE GROUP

Let $V = \bigoplus_{i=0}^m V(\lambda_i)$ ($\lambda_i \in \mathfrak{X}^+$) be a faithful representation of G . Then $G \hookrightarrow \mathbb{V} = \bigoplus \mathrm{End}(V(\lambda_i)) \subseteq \mathrm{End}(V) \cong V \otimes V^*$. Passing to the projectivization, we have $G \hookrightarrow \mathbb{P} = \mathbb{P}(\mathbb{V}) \iff \mathfrak{X} = \sum \mathbb{Z}(\lambda_i - \lambda_j) + Q$. This condition can always be achieved by adding to V a trivial representation, and we will assume it or, more generally, we will consider faithful projective representations $G : \mathbb{P}(V)$ (which can be lifted to a linear representation of rather a finite cover of G , than G itself). Moreover, we may assume all λ_i to be distinct.

Our aim is to study $X = \overline{G} \subseteq \mathbb{P}$.

The image of the identity in $\mathbb{V}$ is $e = \sum v_\mu \otimes v_{-\mu}$, where $v_{\pm\mu}$ run over dual weight bases in V, V^* . Under the above identification $\mathfrak{X}((T \times T)/\mathrm{diag} T) \cong \mathfrak{X}(T)$, we have $\mathrm{supp} e = -\mathcal{P}$.

All closed $(G \times G)$ -orbits in $\mathbb{P}$ are of the form $Y_i = (G \times G)y_i$, $y_i = \langle v_{\lambda_i} \otimes v_{-\lambda_i} \rangle$.

The local structure of $\mathbb{P}$ in a neighborhood of Y_i looks as follows (§6). Let $P = P(\lambda_i)$ be the projective stabilizer of v_{λ_i} , and $P = P_u \rtimes L$, $P^- = P_u^- \rtimes L$ be the Levi decompositions. Then $P \times P^- = (G \times G)_{y_i}$. Consider the open subset

$$\mathring{\mathbb{V}} = \mathbb{C}^\times(v_{\lambda_i} \otimes v_{-\lambda_i}) \oplus (\mathfrak{p}_u^- v_{\lambda_i} \otimes v_{-\lambda_i}) \oplus (v_{\lambda_i} \otimes \mathfrak{p}_u v_{-\lambda_i}) \oplus \mathbb{M}$$

in $\mathbb{V}$, where $\mathbb{M} \subset \mathbb{V}$ is an $(L \times L)$ -stable subspace and $\mathbb{C}(v_{\lambda_i} \otimes v_{-\lambda_i}) \oplus \mathbb{M} \ni e$. The affine $(P^- \times P)$ -stable chart $\mathring{\mathbb{P}} = \mathbb{P}(\mathring{\mathbb{V}})$ intersects Y_i , so that $\mathring{\mathbb{P}} \cong (P^- \times P) \times_{(L \times L)} \mathbb{M}(-\lambda_i, \lambda_i) = P_u^- \times P_u \times \mathbb{M}(-\lambda_i, \lambda_i)$

Proposition 5. $X \supset Y_i \iff \lambda_i$ is a vertex of $\mathcal{P}$

Proof. If λ_i is a vertex of $\mathcal{P}$, then there exists a 1-parameter subgroup $\gamma(t) \in T$ such that $\langle \gamma, \lambda_i \rangle > \langle \gamma, \mu \rangle$, Then

$$\gamma(t) = \left\langle \sum \gamma(t) v_\mu \otimes v_{-\mu} \right\rangle = \left\langle \sum t^{\langle \gamma, \mu \rangle} v_\mu \otimes v_{-\mu} \right\rangle \rightarrow y_i \quad \text{as } t \rightarrow \infty$$

Conversely, assume $Y_i \subset X$. Consider the local structure of X in a neighborhood of Y_i . X contains the $(P^- \times P)$ -stable affine open subset $\mathring{X} = P_u^- \times P_u \times Z$, where $Z = X \cap \mathbb{M}(-\lambda_i, \lambda_i) = \overline{(L \times L)e} = \overline{L}$ is an affine $(L \times L)$ -equivariant embedding of L with the unique fixed point y_i . Then by Theorem 4, $\mathbb{C}[\tilde{Z}] = \bigoplus_{\lambda \in \mathfrak{X} \cap \mathcal{C}^\vee} \mathbb{C}[L]_{(\lambda)}$, where $\mathcal{C}$ is a strictly convex solid cone generated by Δ_L^+ and by finitely many vectors from $\mathcal{V}_L = -C_L$, so that $\text{int } \mathcal{C} \cap \mathcal{V}_L \neq \emptyset$. Hence $\mathbb{C}[Z] = \bigoplus_{\lambda \in S} \mathbb{C}[L]_{(\lambda)}$, where $S \subseteq \mathfrak{X}_L^+ \cap \mathcal{C}^\vee$ is a finitely generated semigroup such that $\mathbb{Z}S \cap \mathcal{C}^\vee = \mathfrak{X} \cap \mathcal{C}^\vee$.

There exists a 1-parameter subgroup $\gamma \in \text{int } \mathcal{C} \cap -C_L$, $\gamma \perp \Delta_L$. It defines a non-negative grading of $\mathbb{C}[Z]$: $\deg \mathbb{C}[Z]_{(\lambda)} = \langle \gamma, \lambda \rangle$, and functions of zero degree are constant. Consequently the action of $\gamma(t)$ contracts the whole Z (and in particular, e) to y_i as $t \rightarrow \infty$, whence $-\lambda_i$ is a vertex of $\text{supp } e = -\mathcal{P}$. $\square$

Let us give a more detailed study of the local structure of X in a neighborhood of a closed orbit, say Y_0 . We keep the previous notation. Put $\mathring{X} = X \cap \mathring{\mathbb{P}} = \{f_{\lambda_0} \neq 0\}$. (Here $f_{\lambda_0} = v_{-\lambda_0} \otimes v_{\lambda_0}$ is regarded as a linear function on $\mathbb{V}$.) Then $\mathring{X} \cong (P^- \times P) \times_{(L \times L)} Z = P_u^- \times P_u \times Z$, where $Z = X \cap (\{y_0\} \times \mathbb{M}) = \overline{L}$ is an equivariant affine embedding.

More precisely, if $V = \mathbb{C}v_{\lambda_0} \oplus V_0$ is an L -stable decomposition, then $Z \subseteq \text{End}(V_0(-\lambda_0)) \subseteq \mathbb{M}(-\lambda_0, \lambda_0)$, whence

Proposition 6. Let $\mu_1, \dots, \mu_s \in \mathfrak{X}_L^+$ be all highest weights of $L : V$, except λ_0 . The representations $L : V_L(\mu_j - \lambda_0)$ extend to Z and induce a closed embedding $Z \hookrightarrow \mathbb{V}_0 = \bigoplus \text{End}(V_L(\mu_j - \lambda_0))$.

Proposition 7. $\mathbb{C}[Z] = \bigoplus_{\lambda \in S} \mathbb{C}[L]_{(\lambda)}$, where $S \subseteq \mathfrak{X}_L^+$ is a semigroup generating the cone Σ_0 of $C \cap \mathcal{P}$ at λ_0 , and $\mathbb{Z}S = \mathfrak{X}$.

Proof. The homogeneous coordinate algebra of $\mathbb{P}$ is $\mathbb{C}[\mathbb{V}] = \mathbf{S}^\bullet \mathbb{V}^*$. The set of weights of homogeneous polynomials of degree n is contained in $-n\mathcal{P} \oplus n\mathcal{P}$. As $\mathbb{C}[\mathring{\mathbb{P}}] = \bigoplus \mathbf{S}^n \mathbb{V}^* / f_{\lambda_0}^n$, the weights of homogeneous polynomials of degree n on $\mathring{\mathbb{P}}$ lie in $n(-\mathcal{P} + \lambda_0) \oplus n(\mathcal{P} - \lambda_0)$. Hence the highest weights of the $(L \times L)$ -module $\mathbb{C}[Z] \subseteq \mathbb{C}[L]$ lie in $C_L \cap \bigcup n(\mathcal{P} - \lambda_0) = \Sigma_0$ (under the antidiagonal embedding $\mathfrak{X} \subset \mathfrak{X}(T \times T)$).

On the other hand, for each $\mu \in C_L \cap \mathcal{P}$, Lemma 1 yields $V^{\otimes n} \hookrightarrow V_L(n\mu)$ for some n . Using Proposition 4 for L , it is easy to derive $f_{n\mu}/f_{\lambda_0}^n \in \mathbb{C}[Z]$. Therefore S generates Σ_0 . $\square$

Corollary. $\mathbb{C}[\tilde{Z}] = \bigoplus_{\lambda \in \mathfrak{X} \cap \Sigma_0} \mathbb{C}[L]_{(\lambda)}$

Remark. If λ_0 is a regular weight (i.e., $\lambda_0 \in \text{int } C$), then $L = T$, Σ_0 is the cone of $\mathcal{P}$ at λ_0 , and $\mathbb{C}[Z] = \mathbb{C}[S]$ is the semigroup algebra of the semigroup S generated by $\mu - \lambda_0$, where μ runs over all weights of V (cf. §5). In the general case, the description of S requires decomposing the G -module V into simple L -modules and also tensor products of simple L -modules (see §10). Normality of Z means $S = \mathfrak{X} \cap \Sigma_0$.

Consider the spherical variety $\tilde{X}$. By Proposition 1, the normalization map $\tilde{X} \rightarrow X$ is bijective on the set of orbits. In particular, $\tilde{X} \hookrightarrow G/H, Y_i$.

Theorem 7. *The colored fan of $\tilde{X}$ is generated by colored cones of closed orbits $Y_i \subset \tilde{X}$ over all vertices $\lambda_i \in \mathcal{P}$: $\mathcal{C}_{Y_i} = \Sigma_i^\vee$, where Σ_i is the cone of $C \cap \mathcal{P}$ at λ_i , and $\mathcal{F}_{Y_i} = \Pi_{L_i}^\vee$ is the subsystem of simple coroots of the respective Levi subgroup $L_i = Z_G(\lambda_i)$. (Recall that colors are identified with simple coroots.)*

Proof. The assertion on $\mathcal{C}_{Y_i}$ stems from Proposition 7 and from arguments in §6. Let us deal with colors.

Note that $D_j \in \mathcal{F}_{Y_i} \iff y_i \in \overline{D_j}$. As $f_{\lambda_i}(y_i) \neq 0$ and f_{λ_i} has order $\langle \lambda_i, \alpha_j^\vee \rangle$ along D_j , it follows that $D_j \notin \mathcal{F}_{Y_i}$ whenever $\lambda_i \not\prec \alpha_j \iff \alpha_j \notin \Pi_{L_i}$. Conversely, assume $\alpha_j \in \Pi_{L_i}$, and let $\gamma(t) \in T$ be a 1-parameter subgroup such that $\langle \gamma, \lambda_i \rangle > \langle \gamma, \mu \rangle$, $\forall \mu \in \mathcal{P}$, $\mu \neq \lambda_i$. Then as $t \rightarrow \infty$,

$$\gamma(t)s_{\alpha_j} = \left\langle \sum \gamma(t)v_{s_{\alpha_j}\mu} \otimes v_{-\mu} \right\rangle = \left\langle \sum t^{\langle \gamma, s_{\alpha_j}\mu \rangle} v_{s_{\alpha_j}\mu} \otimes v_{-\mu} \right\rangle \rightarrow y_i,$$

i.e., $\overline{D_j} \ni y_i$. $\square$

Remark. In case, where all vertices of $\mathcal{P}$ are regular weights, the fan of X has no colors and is a subdivision of $-C$. Transversal slices Z_i to closed orbits Y_i at y_i are affine toric varieties, and their cones are dual to the cones of $\mathcal{P}$ at λ_i . These slices are contained in a projective toric variety $\overline{T} \subseteq X$ (or $\tilde{X}$). Clearly, $\overline{T}$ is W -stable (W or, more precisely, $N_G(T)$ acts by conjugation). It follows from the local structure that $\overline{T}$ intersects all Y_i transversally at the points $w(y_i)$, $w \in W$, which are all T -fixed points of $\overline{T}$. The fan of $\overline{T}$ is the W -span of the fan of X , and its polytope equals $\mathcal{P}$ by Proposition 5.

Left cosets $\overline{vT}$, $v \in W$, intersect all Y_i transversally at the points $(vw, w)y_i$, $w \in W$, which are all $(T \times T)$ -fixed points of X . The closure of the normalizer of the torus $\overline{N_G(T)} = \bigsqcup_{v \in W} \overline{vT}$ is a disjoint union of cosets. (Otherwise distinct $\overline{vT}$ would share fixed points.)

In case, where $V = V(\lambda_0)$ is irreducible, X is normal (even smooth, cf. §§10,11 and [CP-I]) and does not depend on the choice of (regular) λ_0 . This compactification is called *wonderful* [CP-I]. The general case of regular weights was considered in [CP-II] and in a number of other papers.

9. ORBITS

According to the general theory of spherical varieties and Theorem 7, $(G \times G)$ -orbits in X whose closures contain a given closed orbit Y_i are in bijection with colored faces of the colored cone $(\Sigma_i^\vee, \Pi_{L_i}^\vee)$ or, equivalently, with faces $\mathcal{C} \subseteq \Sigma_i^\vee$ such that $(\text{int } \mathcal{C}) \cap (-C) \neq \emptyset$. We reformulate this description in “dual” terms of $\mathcal{P}$.

For any face $\Gamma \subseteq \mathcal{P}$, denote by $|\Gamma|$ its direction subspace, by $\langle \Gamma \rangle$ its linear span, and put $\|\Gamma\| = |\Gamma| \oplus (\langle \Gamma \rangle^\perp \cap \langle Q \rangle)$. (If G is semisimple and Γ is a proper face, then $\|\Gamma\|$ is its supporting hyperplane shifted to 0.) We say that $\gamma \in \text{Hom}(\mathfrak{X}, \mathbb{Q})$ is a *supporting function* for Γ in $\mathcal{P}$ if $\langle \gamma, \Gamma \rangle = \text{const} < \langle \gamma, \mathcal{P} \setminus \Gamma \rangle$, i.e., γ is the l.h.s. of the equation of a supporting hyperplane for Γ .

Proposition 8. *There is a bijection between all $(G \times G)$ -orbits $Y \subset X$ and all faces $\Gamma \subseteq \mathcal{P}$ such that $(\text{int } \Gamma) \cap C \neq \emptyset$. Here $\mathcal{C}_Y$ is dual to the cone of $C \cap \mathcal{P}$ at the face $C \cap \Gamma$, and $\mathcal{F}_Y = \{D_j \mid \alpha_j \perp \langle \Gamma \rangle\}$. “Colored” orbits (i.e., those having $\mathcal{F}_Y \neq \emptyset$) correspond to faces lying at the boundary of C . The adherence of orbits (i.e., inclusion of orbit closures) corresponds to the inclusion of the respective faces.*

Proof. Assume $\overline{Y} \supseteq Y_i$, i.e., $\mathcal{C}_Y$ is a face of $\Sigma_i^\vee$. We have $\gamma \in \text{int } \mathcal{C}_Y \iff \gamma$ is a supporting function for the dual face $\Sigma_Y = \Sigma_i \cap \mathcal{C}_Y^\perp$ of Σ_i , i.e., $\langle \gamma, \Sigma_i \rangle \geq 0$ and Σ_Y is distinguished by the equation $\langle \gamma, \cdot \rangle = 0$. Thus colored faces $\Sigma_i^\vee$ correspond to faces of Σ_i , or of $C \cap \mathcal{P}$ at λ_i , cut out by supporting functions $\gamma \in -C$. But such faces of $C \cap \mathcal{P}$ are obtained by intersecting C with faces $\Gamma \subseteq \mathcal{P}$ of the same dimension cut out by γ in $\mathcal{P}$.

The description of colors stems from $\Pi_{L_i}^\vee = \Pi^\vee \cap \lambda_i^\perp$ and $\mathcal{F}_Y = \Pi_{L_i}^\vee \cap |\Gamma|^\perp$. The assertion on adherence follows from Theorem 3, because duality reverts inclusion of faces. $\square$

Let us make the above description of orbits more explicit. We introduce the following notation.

For any subspace $N \subseteq \mathfrak{X} \otimes \mathbb{Q}$ orthogonal to some dominant weight, denote by P_N the parabolic in G generated by B and by the roots $\alpha \in N$. Let $P_N^\pm = (P_N^\pm)_u \rtimes L_N$ be the Levi decompositions, $L_N \supseteq T$, L'_N be the commutator subgroup of L_N . For any sublattice $\Lambda \subseteq \mathfrak{X}$, denote by $T^\Lambda \subseteq T$ the diagonalizable group which is the common kernel of all characters $\lambda \in \Lambda$.

For any face $\Gamma \subseteq \mathcal{P}$, denote by V_Γ the sum of weight subspaces of V with weights in Γ , and by V'_Γ its T -stable complement. Let $e_\Gamma =$

$\sum_{\mu \in \Gamma} v_\mu \otimes v_{-\mu}$ be the projector on V_Γ w.r.t. the decomposition $V = V_\Gamma \oplus V'_\Gamma$.

Observe that V_Γ, V'_Γ are $L_{|\Gamma|}$ - and even $L_{\|\Gamma\|}$ -stable, $L'_{\|\Gamma\|} = L'_{|\Gamma|} \times L'_{\langle \Gamma \rangle^\perp}$, and the action $L'_{\langle \Gamma \rangle^\perp} : V_\Gamma$ is trivial. Indeed, adding roots $\alpha \in \|\Gamma\| \setminus |\Gamma|$ moves Γ outside $\mathcal{P}$, i.e., the respective root vectors act on V_Γ trivially. This means that $\Delta \cap \|\Gamma\| = (\Delta \cap |\Gamma|) \sqcup (\Delta \cap \langle \Gamma \rangle^\perp)$ is a disjoint orthogonal union.

In the subspaces $|\Gamma|, \langle \Gamma \rangle$, consider generating sublattices

$$\begin{aligned} |\Gamma|_{\mathbb{Z}} &= \sum_{\lambda_i, \lambda_j \in \Gamma} \mathbb{Z}(\lambda_i - \lambda_j) + Q \cap |\Gamma| \\ \langle \Gamma \rangle_{\mathbb{Z}} &= \sum_{\lambda_i \in \Gamma} \mathbb{Z}\lambda_i + Q \cap |\Gamma| \end{aligned}$$

(To be rigorous, these lattices depend not only on $\mathcal{P}$ and Γ , but on the initial set of highest weights $\lambda_0, \dots, \lambda_m$.) Note that $\langle \Gamma \rangle_{\mathbb{Z}}$ is the weight lattice of $T : V_\Gamma$.

Theorem 8. *The orbit $Y \subset X$ corresponding to the face $\Gamma \subseteq \mathcal{P}$ is represented by $y = \langle e_\Gamma \rangle$. Stabilizers look like*

$$\begin{aligned} (G \times G)_y &= ((P_{\|\Gamma\|})_{\mathbf{u}} \times (P_{\|\Gamma\|}^-)_{\mathbf{u}}) \ltimes ((L'_{\langle \Gamma \rangle^\perp} T^{|\Gamma|_{\mathbb{Z}}} \times L'_{\langle \Gamma \rangle^\perp} T^{|\Gamma|_{\mathbb{Z}}}) \cdot \text{diag } L_{|\Gamma|}) \\ \dim Y &= \dim G - \dim L_{\langle \Gamma \rangle^\perp} + \dim \Gamma \end{aligned}$$

Proof. Take $\gamma \in (\text{int } \mathcal{C}_Y) \cap \mathcal{V}$. Then

$$\gamma(t) = \left\langle \sum \gamma(t) v_\mu \otimes v_{-\mu} \right\rangle = \left\langle \sum t^{\langle \gamma, \mu \rangle} v_\mu \otimes v_{-\mu} \right\rangle \rightarrow y \quad \text{as } t \rightarrow 0$$

whence $y \in \overline{T}$. Moreover, $f_\lambda(\gamma(t)) = t^{\langle \lambda, \gamma \rangle} c_\lambda$, $c_\lambda = f_\lambda(e) \neq 0$, $\forall \lambda \in \mathfrak{X}$.

Choose a closed orbit $Y_0 \subseteq \overline{Y}$ corresponding to a certain vertex $\lambda_0 \in C$ of Γ . In the notation of §8, $y \in \dot{X}$, whence $y \in Z$. We have to prove that $y \in Y \cap Z$.

It follows from the local structure of $\dot{X}$ (§6) that $Y \cap Z$ is an $(L \times L)$ -orbit in Z and $\mathcal{C}_{Y \cap Z} = \mathcal{C}_Y$ is a face of $\Sigma_0^\vee$. The ideal of $\overline{Y \cap Z}$ (the closure in Z) as an $(L \times L)$ -submodule in $\mathbb{C}[Z]$ is given by its highest weight vectors f_λ , $\lambda \in \Sigma_0 \setminus \mathcal{C}_Y^\perp = \Sigma_0 \setminus |\Gamma|$. Thence $f_\lambda(y) = 0 \iff f_\lambda|_{Y \cap Z} = 0$.

Furthermore, for all $(b, b^-) \in B \times B^-$ we have

$$(b, b^-) f_\lambda(y) = f_\lambda(\langle b^{-1} e_\Gamma b^- \rangle) = f_\lambda(b_\Gamma^{-1} e_\Gamma b_\Gamma^-)$$

(Here $b^\pm = u_\Gamma^\pm b_\Gamma^\pm$ are the decompositions in $B^\pm = (P_{|\Gamma|}^\pm)_{\mathbf{u}} \ltimes B_{|\Gamma|}^\pm$, $B_{|\Gamma|}^\pm = B^\pm \cap L_{|\Gamma|}$.) The latter expression may be regarded as the value of a function of weight $(-\lambda, \lambda)$ at $b_\Gamma^{-1} b_\Gamma^- \in L_{|\Gamma|}$, since the identity maps to e_Γ under the representation $L_{|\Gamma|} : V_\Gamma$. As the two “big cells” $B_{|\Gamma|} B_{|\Gamma|}^-, B_{|\Gamma|}^- B_{|\Gamma|} \subseteq L_{|\Gamma|}$ intersect in a dense subset, for almost all (b, b^-) there exist $(c_\Gamma^-, c_\Gamma) \in B_{|\Gamma|}^- \times B_{|\Gamma|}$ such that $f_\lambda(b_\Gamma^{-1} e_\Gamma b_\Gamma^-) = f_\lambda(c_\Gamma^- e_\Gamma c_\Gamma) = d_\lambda f_\lambda(y)$, $d_\lambda \neq 0$. (Recall that f_λ is an eigenfunction for $B_{|\Gamma|}^- \times B_{|\Gamma|}$.)

It follows that $f_\lambda(y) = 0$ iff $\mathbb{C}[Z]_{(\lambda)}$ vanishes at y , whence $y \in \overline{Y \cap Z}$ and is not contained in smaller orbits. This implies $y \in Y \cap Z$.

Now we pass to stabilizers. For $(g, h) \in G \times G$ we have: $(g, h)y = y$ iff $ge_\Gamma h^{-1}$ is proportional to e_Γ iff

- (1) $gV_\Gamma = V_\Gamma$,
- (2) $hV'_\Gamma = V'_\Gamma$,
- (3) the actions of g, h on $V_\Gamma \cong V/V'_\Gamma$ differ by a scalar multiple.

The condition (1) means that $g \in P_{\|\Gamma\|}$, (2) $\iff h \in P_{\|\Gamma\|}^-$, and the kernels of the actions $P_{\|\Gamma\|}^\pm : V_\Gamma \cong V/V'_\Gamma$ are $(P_{\|\Gamma\|}^\pm)_u \times (L'_{\langle \Gamma \rangle^\perp} T^{\langle \Gamma \rangle^\mathbb{Z}})$, i.e., we may assume below that $g, h \in L_{|\Gamma|}$. But now (3) just means that, up to multiplying by one and the same element of $L_{|\Gamma|}$, $g, h \in T^{|\Gamma|\mathbb{Z}}$.

The formula for $\dim Y$ easily follows from the structure of $(G \times G)_y$ and of $L'_{\|\Gamma\|}$. $\square$

Remark. The theorem generalizes the results of de Concini–Procesi for wonderful and regular completions (see [CP-I], [CP-II], [Br4]) and those of Putcha–Renner [PR], [Pu2], [Re2], and Vinberg [Vin, Thm.7] for algebraic semigroups. A direct link to algebraic semigroups is provided by considering the cone over X in $\mathbb{V}$. It is an algebraic semigroup, whose group of invertibles is the extension of G by homotheties (cf. [Re2]). The idea of computing stabilizers is taken from [Vin, §7].

The results close to Proposition 8 and Theorem 8 were obtained for *normal* affine and projective embeddings of G (and even in a more general context) by Alexeev and Brion [AB1], [AB2]. When the preliminary text of this paper was written, the author knew about the paper of Kapranov [Kap] in which all assertions of Proposition 8 and Theorem 8, except for stabilizers and dimensions, were proved, see [Kap, 2.4.2]. However the proof therein seems to be incomplete. The author is indebted to M. Brion for this reference.

Corollary. $\overline{T}$ intersects all $(G \times G)$ -orbits in X . T -orbits in the intersection of $\overline{T}$ with a $(G \times G)$ -orbit are permuted by W transitively.

Remark. An assertion similar to the first one holds for any spherical variety [Br3, 2.4]. A simplest “transcendental” proof of the corollary is obtained by closing in X the Cartan decomposition $G = KTK$, where $K \subset G$ is a maximal compact subgroup. One can give an “algebraic” proof in the same way by considering the Iwahori decomposition of $G(\mathbb{C}((t)))$ instead of the Cartan decomposition, cf. [Br3, 2.4, Example 2].

On the other hand, this corollary can be used to obtain a simple proof of Theorem 8 as follows. It is an easy exercise in toric geometry (cf. §5) that $y = \langle e_\Gamma \rangle$ over *all* faces $\Gamma \subseteq \mathcal{P}$ form a complete set of T -orbit representatives in $\overline{T}$. Thus these y represent all $(G \times G)$ -orbits in X . Now it is easy to deduce from the structure of the respective orbit Y as a homogeneous space given by $(G \times G)_y$ that $Y^{\text{diag } T}$ is a

union of T -orbits permuted by W transitively. Therefore $y = \langle e_\Gamma \rangle$ over those $\Gamma \subseteq \mathcal{P}$ with $(\text{int } \Gamma) \cap C \neq \emptyset$ form a complete set of $(G \times G)$ -orbit representatives in X .

However we are also interested in the combinatorial data of the embedding theory (colored cones) related to these orbits, so our proof is different.

10. NORMALITY

The questions of normality and smoothness are of local nature. Thus it suffices to examine them at points of some closed orbit $Y_0 \subset X$. A general normality criterion is essentially contained in Propositions 6, 7 and the subsequent remarks.

We say that weights $\mu_1, \dots, \mu_s \in \mathfrak{X}^+$ G -generate a semigroup S if S consists of all highest weights $k_1\mu_1 + \dots + k_s\mu_s - \beta$ ($\beta \in \mathbb{Z}_+\Pi$) of G -modules $V(\mu_1)^{\otimes k_1} \otimes \dots \otimes V(\mu_s)^{\otimes k_s}$, $k_1, \dots, k_s \in \mathbb{Z}_+$.

Proposition 9. *In the notation of §8, let $\mu_1, \dots, \mu_s \in \mathfrak{X}_L^+$ be all highest weights of $L : V$, except λ_0 . Then X is normal at points of Y_0 iff $\mu_1 - \lambda_0, \dots, \mu_s - \lambda_0$ L -generate $\mathfrak{X} \cap \Sigma_0$.*

Proof. Note that X is normal at points of Y_0 iff Z is normal. By Proposition 6, $Z \hookrightarrow \mathbb{V}_0 = \bigoplus \text{End}(V_L(\mu_j - \lambda_0))$, hence $\mathbb{C}[Z] \subseteq \mathbb{C}[L]$ is generated by the components $\mathbb{C}[L]_{(\mu_j - \lambda_0)}$. By Proposition 4, $\mathbb{C}[Z] = \bigoplus_{\lambda \in S} \mathbb{C}[L]_{(\lambda)}$, where $S \subseteq \mathfrak{X}_L^+$ is the semigroup L -generated by $\mu_j - \lambda_0$, $j = 1, \dots, s$. Now the assertion follows from the corollary of Proposition 7. $\square$

In order to apply this criterion, one requires the information on decomposing V into simple L -modules and on decomposing tensor products of L -modules. The first problem is eliminated by the following lemma:

Lemma 3. *The weights $\lambda_1 - \lambda_0, \dots, \lambda_m - \lambda_0$, $-\alpha_j \in -(\Pi \setminus \Pi_L)$ are highest weights of $L : V(-\lambda_0)$ and they L -generate all highest weights of $L : V(-\lambda_0)$.*

Proof. Observe that $v_{\lambda_i}, e_{-\alpha_j}v_{\lambda_0}$ are highest weight vectors of $L : V$, whence the first assertion. Furthermore, for every $i = 0, \dots, m$ there exists a unique L -submodule $V_L(\lambda_i) \hookrightarrow V$ (generated by v_{λ_i}) and

$$V = \sum_{k,i} \underbrace{\mathfrak{g} \cdots \mathfrak{g}}_k v_{\lambda_i} = \sum_{k,i} \underbrace{\mathfrak{p}^- \cdots \mathfrak{p}^-}_k \cdot V_L(\lambda_i)$$

The algebra $\mathfrak{p}^-$ is generated by $\mathfrak{l}$ and $e_{-\alpha_j}$, $\alpha_j \notin \Pi_L$, and it contains simple L -submodules $\mathfrak{g}_L(-\alpha_j) \cong V_L(-\alpha_j)$ with highest weight vectors $e_{-\alpha_j}$. Therefore

$$V = \sum_{n,i,j_1,\dots,j_n} \mathfrak{g}_L(-\alpha_{j_1}) \cdots \mathfrak{g}_L(-\alpha_{j_n}) \cdot V_L(\lambda_i)$$

But $\mathfrak{g}_L(-\alpha_{j_1}) \cdots \mathfrak{g}_L(-\alpha_{j_n}) \cdot V_L(\lambda_i)$ is a quotient module of $V_L(-\alpha_{j_1}) \otimes \cdots \otimes V_L(-\alpha_{j_n}) \otimes V_L(\lambda_i)$. This yields the second assertion of the lemma. $\square$

Thus in Proposition 9 one may replace $\mu_1 - \lambda_0, \dots, \mu_s - \lambda_0$ by $\lambda_1 - \lambda_0, \dots, \lambda_m - \lambda_0, -\alpha_j \in -(\Pi \setminus \Pi_L)$. (Although sometimes it is more convenient to operate with all L -highest weights if they are known.) However the problem of decomposing tensor products of modules with L -generating highest weights into simple L -modules remains. (For normality one has to obtain all generators of $\mathfrak{X} \cap \Sigma_0$ among highest weights of all occurring simple L -submodules.) That is why this criterion is not really effective in the general case. However it implies simple sufficient conditions.

Corollary. *X is normal (at points of Y_0) if $\lambda_1 - \lambda_0, \dots, \lambda_m - \lambda_0$ generate $\mathfrak{X} \cap \Sigma_0$.*


The necessary condition for normality obtained by Renner for reductive algebraic semigroups (see [Re1, 6.4]) extends to projective compactifications.

Proposition 10. *If X is normal, then $\overline{T}$ is normal.*

Proof. Replacing G by L , X by Z , $\overline{T}$ by the closure of T in L , we reduce the problem to the case of an affine embedding. We have $Z \subseteq \mathbb{V}_0 = \bigoplus \text{End}(V_L(\nu_j))$, where $\nu_j \in \mathfrak{X}_L^+$ L -generate $\Sigma_0 \cap \mathfrak{X}$. We can increase $\mathbb{V}_0$ by adding new L -highest weights in such a way that ν_j will generate $\Sigma_0 \cap \mathfrak{X}$. Then $Z = \overline{Z} \subseteq \mathbb{V}_0$ will not change (the highest weights of $\mathbb{C}[Z]$ are the same), but now the weights of $T : V_L(\nu_j)$ will generate the semigroup $W_L \Sigma_0 \cap \mathfrak{X}$ of all lattice points of $W_L \Sigma_0$, the cone of $\mathcal{P}$ at λ_0 . The respective semigroup algebra is the coordinate algebra of $\overline{T} \subseteq Z$, whence $\overline{T}$ is normal. $\square$

Remark. This condition can be effectively verified (§5), and in the case, where all vertices of $\mathcal{P}$ are regular weights, the condition coincides with the general normality criterion. However this condition is not sufficient in the general case.

Example. Let $G = \text{Sp}_4$ and the highest weights of V be $\{\lambda_0, \lambda_1\} = \{3\omega_1, 2\omega_2\}$. The weight polytope $\mathcal{P}$ is given at Fig. (a), the highest weights are indicated by bold dots.



Here $L = \mathrm{SL}_2 \times \mathbb{C}^*$, $\Delta_L = \{\pm\alpha_2\}$. The weight semigroup of $\mathbb{C}[\overline{T}]$ (the closure is taken in Z) is indicated by dots at Fig. (b). Bold dots indicate the subsemigroup of highest weights of $\mathbb{C}[Z]$ (which is easy to compute using the Clebsch–Gordan formula). Now we can see that $\overline{T}$ is normal and Z is not.

11. SMOOTHNESS

In the theory of spherical varieties, smoothness is usually a much more subtle property than normality. The general smoothness criterion [Br2, 4.2] is rather intricate. Surprisingly, for projective compactifications of reductive groups, it is easier to verify smoothness, than normality. We retain the notation of §§8,10.

Theorem 9. *X is smooth at points of Y_0 iff $L \cong \mathrm{GL}_{n_1} \times \cdots \times \mathrm{GL}_{n_p}$ is a direct product, and only polynomial representations of L occur in the decomposition of the L -module $V_0(-\lambda_0)$ (here $V = \mathbb{C}v_{\lambda_0} \oplus V_0$ as above), and all minimal representations of factors occur among them. In the language of roots and weights, this amounts to the following conditions:*

- (1) *All simple components of L are of type A, and there are no more than $\dim Z(L)$ of them.*
- (2) *Σ_0 is simplicial, and moreover, is generated by a basis of $\mathfrak{X}$.*
- (3) *One can enumerate the simple roots in order of their positions at Dynkin diagrams of connected components $\{\alpha_1^{(k)}, \dots, \alpha_{n_k-1}^{(k)}\}$ of Π_L , $k = 1, \dots, q$, and partition the basis of the free semigroup $\mathfrak{X} \cap \Sigma_0$ into subsets $\{\pi_1^{(k)}, \dots, \pi_{n_k}^{(k)}\}$, $k = 1, \dots, p$, $p \geq q$, in such a way that $\langle \pi_j^{(k)}, \alpha_j^{(k)\vee} \rangle = 1$, $\pi_j^{(k)} \perp \Pi_L \setminus \{\alpha_j^{(k)}\}$, $n_k \pi_j^{(k)} - j \pi_{n_k}^{(k)} \perp \pi_{n_1}^{(1)}, \dots, \pi_{n_p}^{(p)}$, $\forall j, k$.*
- (4) *Among the weights $\lambda_1 - \lambda_0, \dots, \lambda_m - \lambda_0$, $-\alpha_j \in -(\Pi \setminus \Pi_L)$ there occur all $\pi_1^{(k)}$, $1 \leq k \leq p$.*

Proof. Observe that X is smooth (at points of Y_0) iff Z is smooth, i.e., the problem is reduced to affine embeddings.

As above, let $\mu_1, \dots, \mu_s$ be the highest weights of $L : V$. We have an embedding $Z \hookrightarrow \mathbb{V}_0 = \bigoplus \mathrm{End}(V_L(\mu_j - \lambda_0))$, $y_0 \mapsto 0$.

If Z is smooth, then by Luna's fundamental lemma (§3), Z projects onto T_0Z isomorphically under an $(L \times L)$ -equivariant projection $\mathbb{V}_0 \rightarrow T_0Z$. Renumeration of μ_j yields

$$Z \cong \text{End}(V_L(\mu_1 - \lambda_0)) \oplus \cdots \oplus \text{End}(V_L(\mu_p - \lambda_0)), \quad p \leq s$$

Let $e \mapsto (e_1, \dots, e_p)$ under this isomorphism. The projection $Z \rightarrow \text{End}(V_L(\mu_k - \lambda_0))$ maps the dense orbit $L \subset Z$ onto the dense orbit Le_k , whence e_k is a nonzero scalar operator. After rescaling the above isomorphism, we may assume e_k to be the identity operator. Then the projection maps L homomorphically onto $\text{GL}(V_L(\mu_k - \lambda_0))$. By a dimension argument, $L \cong \text{GL}_{n_1} \times \cdots \times \text{GL}_{n_p}$, $n_k = \dim V_L(\mu_k - \lambda_0)$, and $\mu_k - \lambda_0$ are the highest weights of minimal representations of GL_{n_k} , $k = 1, \dots, p$.

The semigroup $\mathfrak{X} \cap \Sigma_0$ is L -generated by all $\mu_k - \lambda_0$ and consists of highest weights of all polynomial representations of L (i.e., those extendible to Z). The description of polynomial representations of GL_n implies that $\mathfrak{X} \cap \Sigma_0$ is freely generated by the weights $\pi_j^{(k)}$, where $\pi_1^{(k)}$ is the highest weight of a minimal representation of GL_{n_k} and $V_L(\pi_j^{(k)}) = \bigwedge^j V_L(\pi_1^{(k)})$. Conditions (1)–(4) are easily deduced in view of Lemma 3.

Conversely, if conditions (1)–(4) are satisfied, then $L \cong \text{GL}_{n_1} \times \cdots \times \text{GL}_{n_p}$, and highest weights in $\mathbb{C}[Z]$ correspond to all polynomial representations of L . Therefore $Z \cong \text{Mat}_{n_1} \times \cdots \times \text{Mat}_{n_p}$, i.e., it is smooth. $\square$

Remark. In the case of a regular weight λ_0 , $L = T$, and the smoothness criterion is reduced to conditions (2) and (4): X is smooth (at points of Y_0) $\iff \Sigma_0$ is generated by the basis $\pi^{(1)}, \dots, \pi^{(p)}$ of $\mathfrak{X}$ and $\lambda_0 + \pi^{(k)}$ are in the weight system of $T : V$.

Example. Let $G = \text{SO}_{2l+1}$ and $V = V(\lambda_0)$ be an irreducible representation of the fundamental highest weight $\lambda_0 = \omega_i$. For $i < l$, X will be singular, because it violates condition (1) (or (3) for $l = 2$).

In case of the spinor representation, we have $\lambda_0 = (\varepsilon_1 + \cdots + \varepsilon_l)/2$, where $\pm\varepsilon_1, \dots, \pm\varepsilon_l$ are the nonzero weights of the tautological representation of SO_{2l+1} . The weights of the spinor representation are $(\pm\varepsilon_1 \pm \cdots \pm \varepsilon_l)/2$, $\Pi = \{\varepsilon_1 - \varepsilon_2, \dots, \varepsilon_{l-1} - \varepsilon_l, \varepsilon_l\}$, $\Pi_L = \Pi \setminus \{\varepsilon_l\}$. The highest weights of $L : V$ are $\mu_1 = (\varepsilon_1 + \cdots + \varepsilon_{l-1} - \varepsilon_l)/2$, $\mu_2 = (\varepsilon_1 + \cdots - \varepsilon_{l-1} - \varepsilon_l)/2, \dots, \mu_l = (-\varepsilon_1 - \cdots - \varepsilon_{l-1} - \varepsilon_l)/2$. The vectors $\pi_k = \mu_k - \lambda_0$ generate Σ_0 and form a basis of $\mathfrak{X} = \langle \varepsilon_1, \dots, \varepsilon_l \rangle$. Indeed, $\pi_1 = -\varepsilon_1, \pi_2 = -\varepsilon_1 - \varepsilon_2, \dots, \pi_l = -\varepsilon_1 - \cdots - \varepsilon_l$. It is also easy to see that condition (3) is verified. Thus X is smooth.

Remark. Using a more subtle version of the local structure in the neighborhood of a non-closed orbit, one can obtain conditions of normality and smoothness at any point of X .

12. EXAMPLES

Here we illustrate the general theorems proven above by describing geometric properties of equivariant compactifications of simple algebraic groups in the spaces of projective linear operators of fundamental and adjoint representations. In each case, we describe the structure of the orbit set and examine normality and smoothness of the compactification.

Our results are presented in Tables 1–2. We consider a representation of a simple algebraic group G , indicated up to isomorphism in the column “Group”, in a module $V = V(\lambda_0)$, where λ_0 is a fundamental weight or the highest root. Fundamental representations are denoted in the column “Module” by indicating the highest weight λ_0 and those adjoint representations which are not fundamental are denoted by the symbol Ad. The numeration of simple roots and of fundamental weights, respectively, is taken from [VO] (so that $V(\omega_1)$ always has the minimal dimension).

We consider a $(G \times G)$ -variety $X = \overline{\text{Ad } G} \subseteq \mathbb{P}(\text{End}(V))$. In the column “Orbits” we indicate the dimensions and the Hasse graph of $(G \times G)$ -orbits in X . For classical groups, we also indicate orbit representatives (given by projectors $V \twoheadrightarrow V_\Gamma$; if $V = \mathfrak{g}$, then V_Γ is the center of $(\mathfrak{p}_{|\Gamma|})_{\mathfrak{u}}$ or $(\mathfrak{p}_{\|\Gamma\|})_{\mathfrak{u}}$). These data can be easily derived from Theorem 8 and the following lemma (cf. [PR, Thm. 2]):

Lemma 4. *Let $V = V(\lambda_0)$, $\lambda_0 \in \mathfrak{X}^+$, and $L \subseteq P = P(\lambda_0)$ be the standard Levi subgroup. The faces of $\mathcal{P} = \mathcal{P}(V)$ intersecting C are of the form $\Gamma = \mathcal{P} \cap (\lambda_0 + \langle \Pi_\Gamma \rangle)$, where $\langle \Pi_\Gamma \rangle \subseteq \Pi$ is a subsystem of simple roots such that no connected component of Π_Γ is contained in Π_L , or $\Pi_\Gamma = \emptyset$. Furthermore, $|\Gamma| = \langle \Pi_\Gamma \rangle$, the systems of simple roots of $L_{|\Gamma|}$ and of $L_{\langle \Gamma \rangle^\perp}$ are Π_Γ and $\Pi_{\Gamma^\perp} = \Pi_L \cap (\Pi_\Gamma)^\perp$, respectively, and e_Γ is the $L_{|\Gamma|}$ -equivariant projector onto $V_{L_{|\Gamma|}}(\lambda_0) \subseteq V_G(\lambda_0)$.*

In particular, for $\lambda_0 = \omega_i$, Π_Γ is a connected subsystem of Π containing α_i , or $\Pi_\Gamma = \emptyset$, and $\Pi_{\Gamma^\perp} = (\Pi_\Gamma)^\perp \cap \Pi \setminus \{\alpha_i\}$.

Proof. The faces $\Gamma \subseteq \mathcal{P}$, $\text{int } \Gamma \cap C \neq \emptyset$, are cut out by supporting functions $\gamma \in -C$. Consider the root subsystem $\Delta_\gamma = \Delta \cap \gamma^\perp$ with simple roots in $\Pi_\gamma = \Delta_\gamma \cap \Pi$. The space $|\Gamma|$ is generated by those $\alpha \in \Delta_\gamma^+$ such that $e_{-\alpha} v_{\lambda_0} \neq 0$, i.e., $\alpha \in \Delta_\gamma^+ \setminus \Delta_L^+$. In other words, α has a strictly positive linear expression in a connected subsystem $\Pi' \subseteq \Pi_\gamma$, $\Pi' \not\subseteq \Pi_L$. All such Π' are linearly expressed in such α . Hence $|\Gamma| = \langle \Pi_\Gamma \rangle$, where Π_Γ is the union of connected components of Π_γ that are not contained in Π_L . This yields the description of Γ . The description of e_Γ stems from $V_\Gamma = V_{L_{|\Gamma|}}(\lambda_0)$, and other assertions of the lemma are evident. $\square$

The Hasse diagrams of orbit sets coincide with those for irreducible simple algebraic semigroups (which are nothing else, but the cones over our projective compactifications). The latter are computed in [PR].

We indicate the Levi subgroup of $P = P(\lambda_0)$ in the column “ L ” and the nonzero highest weights of $L : V(-\lambda_0)$ in the column “ L -weights”. If there are too many of them, then we indicate only $-\alpha_j \in -(\Pi \setminus \Pi_L)$ (which L -generate all other highest weights by Lemma 3). We use the following notation: ε is a fixed basic weight of the central 1-torus; $\varepsilon_1, \dots, \varepsilon_n$ are the weights of the tautological representation in $\mathbb{C}^n$ of a classical subgroup of GL_n ($\varepsilon_{n-i} = -\varepsilon_i$ for the orthogonal and the symplectic group); $\pi_i = \varepsilon_1 + \dots + \varepsilon_i$ is the highest weight of $\bigwedge^i \mathbb{C}^n$; if L is represented as a quotient of a direct product of several groups, then the weights of the factors are distinguished by superscripts $', ', \dots$.

In the columns “Normality” and “Smoothness” we indicate whether X has the respective property. If normality fails, we give a reason for it (see Propositions 7,9): if S is the semigroup of highest weights of $L : \mathbb{C}[Z]$ and $\mu \notin S$, but $k\mu \in S$, then we write “ $\nexists \mu, \exists k\mu$ ”. The following two lemmas are helpful in verifying normality:

Lemma 5. *Suppose $G_i : V_i$ ($i = 1, \dots, s$) are faithful representations of connected reductive groups such that $Z(G_i)$ act by homotheties. Let Z_i, Z be the closures of the images of $G_i, G_1 \times \dots \times G_s$ in $\mathrm{End}(V_i), \mathrm{End}(V_1 \otimes \dots \otimes V_s)$, respectively. Then Z is normal iff $Z_1, \dots, Z_s$ are normal.*

Proof. The algebra $\mathbb{C}[Z_1 \times \dots \times Z_s]$ is multi-graded by the action of the s -dimensional torus $Z(G_1) \times \dots \times Z(G_s)$, and $\mathbb{C}[Z]$ is the invariant algebra of the subtorus $T_0 = \{(t_1, \dots, t_s) \mid t_1 \dots t_s = 1\}$. Thus normality of Z is implied by normality of $Z_1, \dots, Z_s$, of direct products and of quotients of normal varieties [Kr, App. I, 4.4, II.3.3, Satz 1].

On the other hand, the action $Z(G_i) : Z_i$ by homotheties lifts to $\tilde{Z}_i$. Let $f_i \in \mathbb{C}[\tilde{Z}_i]$ be an arbitrary homogeneous function. For all $j \neq i$ choose homogeneous functions $f_j \in \mathbb{C}[\tilde{Z}_j]$ of the same degree and consider $f = f_1 \dots f_s \in \mathbb{C}[Z_1 \times \dots \times Z_s]^{T_0}$. It is a rational function on $Z_1 \times \dots \times Z_s$ which is constant on T_0 -orbits. It is easy to see that all T_0 -orbits in $(Z_1 \setminus \{0\}) \times \dots \times (Z_s \setminus \{0\})$ are closed in $Z_1 \times \dots \times Z_s$, i.e., generic T_0 -orbits are closed. Since closed orbits are separated by invariant polynomials, f is pulled back from a rational function on Z which is integral over $\mathbb{C}[Z]$ [Kr, App. I, 3.7, Satz 2, III.3.3, Satz 1]. Thus if Z is normal, then $f \in \mathbb{C}[Z]$, whence $f_i \in \mathbb{C}[Z_i]$. Therefore Z_i is normal. $\square$

Lemma 6. *Let G be one of the groups $\mathrm{GL}_n, \mathbb{C}^\times \times \mathrm{Sp}_{2l}, \mathbb{C}^\times \times \mathrm{SO}_{2l}, \mathbb{C}^\times \times \mathrm{Spin}_n, \mathbb{C}^\times \times \mathrm{E}_l$ ($l = 6, 7$), and $V = V(\mu_0)$, $\mu_0 = \pi_k, \varepsilon + \omega_1, \varepsilon + \omega_1, \varepsilon + \omega_{[n/2]}, \varepsilon + \omega_1$, respectively. Then the closure $Z \subseteq \mathrm{End}(V)$ of the image of G is normal.*

Proof. Let S be the semigroup of highest weights of $G : \mathbb{C}[Z]$ (which is G -generated by μ_0). By Proposition 7 and the subsequent remarks, Z is normal iff S is *saturated*, i.e., $S = \mathfrak{X} \cap \Sigma_0$, where $\mathfrak{X} = \mathbb{Z}S$, and $\Sigma_0 = \mathbb{Q}_+ S$ is the cone over $C \cap \mathcal{P}$.

In the case $G = \mathrm{GL}_n$, all weights in S are polynomial, i.e., are of the form $\mu = m_1 \varepsilon_1 + \cdots + m_n \varepsilon_n$, $m_1 \geq \cdots \geq m_n \geq 0$. It easily follows from the Pieri formula for the decomposition of $V(\mu) \otimes V(\pi_k)$, where μ is any polynomial weight, that S is the saturated semigroup consisting of all polynomial weights μ such that $m_2 + \cdots + m_n \geq (k-1)m_1$ and $m_1 + \cdots + m_n$ is divisible by k .

In the case $G = \mathbb{C}^\times \times \mathrm{Sp}_{2l}$, the dominant weights of $V(\mu_0)^{\otimes m}$ are $\mu = m\varepsilon + m_1 \varepsilon_1 + \cdots + m_l \varepsilon_l$, $m_1 \geq \cdots \geq m_l \geq 0$, $m - m_1 - \cdots - m_l \in 2\mathbb{Z}_+$. All of them are in S . Indeed, $k\varepsilon + \varepsilon_1 + \cdots + \varepsilon_k$ are highest weights of $\bigwedge^k V(\mu_0)$, 2ε is a highest weight of $V(\mu_0)^{\otimes 2}$, and all other μ are their $\mathbb{Z}_+$ -linear combinations. Thus S is saturated.

The situation in the case $G = \mathbb{C}^\times \times \mathrm{SO}_{2l}$ is similar, but dominant weights are of the form $\mu = m\varepsilon + m_1 \varepsilon_1 + \cdots + m_{l-1} \varepsilon_{l-1} \pm m_l \varepsilon_l$.

In the case $G = \mathbb{C}^\times \times \mathrm{Spin}_{2l+1}$, dominant weights of $V(\mu_0)^{\otimes m}$ are of the form $\mu = m\varepsilon + (m_1 \varepsilon_1 + \cdots + m_l \varepsilon_l)/2$, $m \geq m_1 \geq \cdots \geq m_l \geq 0$, $m \equiv m_1 \equiv \cdots \equiv m_l \pmod{2}$. All of them are in S . Indeed, $2\varepsilon + \varepsilon_1 + \cdots + \varepsilon_k$ ($0 \leq k < l$) are highest weights of $V(\mu_0)^{\otimes 2}$, $\mu_0 = \varepsilon + (\varepsilon_1 + \cdots + \varepsilon_l)/2$, and all other μ are their $\mathbb{Z}_+$ -linear combinations. Thus S is saturated.

In the case $G = \mathbb{C}^\times \times \mathrm{Spin}_{2l}$, dominant weights of $V(\mu_0)^{\otimes m}$ are of the form $\mu = m\varepsilon + (m_1 \varepsilon_1 + \cdots + m_l \varepsilon_l)/2$, $m \geq m_1 \geq \cdots \geq m_{l-1} \geq |m_l|$, $(l-2)m \geq m_1 + \cdots + m_{l-1} - m_l$, $m \equiv m_1 \equiv \cdots \equiv m_l \pmod{2}$, $lm \equiv m_1 + \cdots + m_l \pmod{4}$. These conditions determine a saturated semigroup, which is generated by $\mu_0 = \varepsilon + \omega_l$, $2\varepsilon + \omega_{l-2k}$ ($1 \leq k \leq l/2$), $4\varepsilon + \omega_i + \omega_j$ ($0 \leq i \leq j \leq l-2$, $i \equiv j \pmod{2}$), $k\varepsilon + \omega_{l-i} + (k-2)\omega_{l-1}$ ($2 \leq k \leq i \leq l$, $k \equiv i \pmod{2}$), $(k+2)\varepsilon + \omega_{l-i} + \omega_{l-j} + (k-2)\omega_{l-1}$ ($2 \leq k \leq i \leq j \leq l$, $i \equiv j+k \pmod{2}$). (Here $\omega_0 = 0$.) It is easy to verify that all of them are highest weights in the respective $V(\mu_0)^{\otimes m}$ (where m is the coefficient in ε). Thus our semigroup coincides with S .

A similar reasoning applies to $G = \mathbb{C}^\times \times \mathrm{E}_l$. One considers a semigroup $\mathfrak{X} \cap \Sigma_0$, where $\Sigma_0 = \mathbb{Q}_+(C \cap \mathcal{P})$. It is generated by the following weights: $k\varepsilon + \omega_k$, $(k+9-l)\varepsilon + \omega_k$ ($0 \leq k \leq l-3$, $\omega_0 = 0$), $(l-2)\varepsilon + \omega_{l-2} + \omega_l$, $(l-1)\varepsilon + 2\omega_{l-2}$, $4\varepsilon + \omega_{l-2}$, $2\varepsilon + \omega_{l-1}$, $(l-1)\varepsilon + \omega_{l-1} + 2\omega_l$, $l\varepsilon + 3\omega_l$, $3\varepsilon + \omega_l$. One verifies that all of them occur as highest weights in $V(\mu_0)^{\otimes m}$, whence $S = \mathfrak{X} \cap \Sigma_0$. We omit routine computations. $\square$

Remark. Observe that the list of μ_0 restricted to the maximal torus of G' in Lemma 6 is nothing else but the list of minuscule weights of simple algebraic groups, up to diagram automorphisms. It is obvious that if (the restriction of) μ_0 is not minuscule, then the semigroup S is not saturated, whence Z is not normal. This was first observed by Faltings [Fal], and he also proved the normality of Z for some minuscule weights. The uniform proof of normality for any minuscule weight was recently

obtained by de Concini [Con] using a representation-theoretic lemma from [Kan]. When the preliminary text of this paper was written, the author learned about the papers [Fal], [Con], [Kan] from M. Brion, to whom he is grateful for these references.

Table 1: Orbits and adherence

Group	Module	Orbits
$A_l = \mathrm{SL}_n$ $n = l + 1$	ω_1, ω_l	$Y_r \ni \langle \phi \rangle, \mathrm{rk} \phi = r, \dim Y_r = 2nr - r^2 - 1$ $Y_n \longrightarrow \cdots \longrightarrow Y_1$
	ω_k $1 < k < l$	$Y_{sr} \ni \langle \bigwedge^k \mathbb{C}^n \twoheadrightarrow \bigwedge^s \mathbb{C}^s \otimes \bigwedge^{k-s} \mathbb{C}^{r-s} \rangle$ $\dim Y_{sr} = 2nr - r^2 - s^2 - 1, \dim Y_{kk} = 2nk - 2k^2$ $Y_{0n} \longrightarrow \cdots \longrightarrow Y_{0,k+1}$ $\downarrow \quad \quad \downarrow \quad \quad \downarrow$ $\vdots \quad \quad \vdots \quad \quad \vdots$ $\downarrow \quad \quad \downarrow \quad \quad \downarrow$ $Y_{k-1,n} \longrightarrow \cdots \longrightarrow Y_{k-1,k+1} \longrightarrow Y_{kk}$
	Ad	$\dim Y_0 = n^2 - 1, \dim Y_{sr} = n^2 - (r-s)^2 - 2$ $Y_0 \begin{matrix} \nearrow Y_{11} \\ \nearrow Y_{22} \\ \vdots \\ \searrow Y_{n-1,n-1} \end{matrix} \begin{matrix} \nearrow Y_{12} \\ \nearrow \vdots \\ \searrow Y_{1,n-1} \\ \searrow Y_{n-2,n-1} \end{matrix}$
$B_l = \mathrm{SO}_n$ $n = 2l + 1$	ω_k $k < l$	$Y_{sr} \ni \langle \bigwedge^k \mathbb{C}^n \twoheadrightarrow \bigwedge^s \mathbb{C}^s \otimes \bigwedge^{k-s} \mathbb{C}^{r-s} \rangle$ $Y_s \ni \langle \bigwedge^k \mathbb{C}^n \twoheadrightarrow \bigwedge^s \mathbb{C}^s \otimes \bigwedge^{k-s} \mathbb{C}^{n-2s} \rangle$ $\mathbb{C}^s \perp \mathbb{C}^{n-2s} \supset \mathbb{C}^{r-s}$ are subspaces in $\mathbb{C}^n$ $\mathbb{C}^s, \mathbb{C}^{r-s}$ are isotropic and $\mathbb{C}^{n-2s}$ is nondegenerate $\dim Y_{sr} = (2n - 2r - 1)r - s^2 - 1$ $\dim Y_{kk} = (2n - 3k - 1)k, \dim Y_s = n(n-1)/2 - s^2$ $Y_0 \longrightarrow Y_{0l} \longrightarrow \cdots \longrightarrow Y_{0,k+1}$ $\downarrow \quad \quad \downarrow \quad \quad \downarrow \quad \quad \downarrow$ $\vdots \quad \quad \vdots \quad \quad \vdots \quad \quad \vdots$ $\downarrow \quad \quad \downarrow \quad \quad \downarrow \quad \quad \downarrow$ $Y_{k-1} \longrightarrow Y_{k-1,l} \longrightarrow \cdots \longrightarrow Y_{k-1,k+1} \longrightarrow Y_{kk}$
	ω_l	$Y_s \ni \langle S^n \twoheadrightarrow \bigwedge^s \mathbb{C}^s \otimes S^{n-2s} \rangle$ S^m is the spinor space over $\mathbb{C}^m$ $\mathbb{C}^s \perp \mathbb{C}^{n-2s}$ are subspaces in $\mathbb{C}^n$ $\mathbb{C}^s$ is isotropic and $\mathbb{C}^{n-2s}$ is nondegenerate $\dim Y_s = n(n-1)/2 - s^2$ $Y_0 \longrightarrow \cdots \longrightarrow Y_l$

Table 1: (continued)







Group	Module	Orbits
$C_l = Sp_n$ $n = 2l$	ω_k	Orbits and their adherence are similar to the case B_l , only $\bigwedge^{k-s} \mathbb{C}^{n-2s}$ ($s = 0, \dots, k$) is replaced with its highest component; $\dim Y_{sr} = (2n - 2r + 1)r - s^2 - 1$, $\dim Y_{kk} = (2n - 3k + 1)k$, $\dim Y_s = n(n + 1)/2 - s^2$
	Ad	$\dim Y = n(n + 1)/2$, $\dim Y_r = (2n - 2r + 1)r - 1$ $Y \longrightarrow Y_l \longrightarrow \dots \longrightarrow Y_1$
$D_l = SO_n$ $n = 2l$	ω_k $k \leq l - 2$	Y_s, Y_{sr} are the same as for B_l , but Y_{sl} splits in 2 orbits $Y_{sl}^\pm$ related to the choice of $\mathbb{C}^l = \mathbb{C}^s \oplus \mathbb{C}^{l-s}$ in one of the two connected components of the isotropic Grassmannian. 
	ω_{l-1}, ω_l	$Y_s \ni \langle S_\pm^n \rightarrow \bigwedge^s \mathbb{C}^s \otimes S_\pm^{n-2s} \rangle$ $Y'_s \ni \langle S_\pm^n \rightarrow \bigwedge^s \mathbb{C}^l \otimes \mathbb{C}^{l-s} \rangle$ $S_\pm^m \subset S^m$ are the semispinor spaces $\mathbb{C}^l = \mathbb{C}^s \oplus \mathbb{C}^{l-s} \subset \mathbb{C}^n$ is isotropic $\dim Y_s = n(n - 1)/2 - s^2$, $\dim Y'_s = n(n - 1)/2 - s^2 - 1$ $(0 \leq s \leq l - 3)$ $\dim Y_{l-2} = l^2 + 3l - 7$, $\dim Y_{l-1} = l(l - 1)$ 
G_2	ω_1, ω_2	$14 \rightarrow 13 \rightarrow 10$
F_4	ω_1, ω_4	$52 \rightarrow 51 \rightarrow 48 \rightarrow 43 \rightarrow 30$
	ω_2, ω_3	
E_6	ω_1, ω_5	
	ω_2, ω_4	
	ω_3	

Table 1: (continued)

Group	Module	Orbits
	ω_6	
E_7	ω_1	
	ω_2	
	ω_3	
	ω_4	
	ω_5	
	ω_6	
	ω_7	

Table 1: (continued)

Group	Module	Orbits
E_8	ω_1	$ \begin{array}{ccccccccccc} 248 & \rightarrow & 247 & \rightarrow & 244 & & & & & & \\ \downarrow & & \downarrow & & \downarrow & & & & & & \\ 247 & \rightarrow & 246 & \rightarrow & 243 & \rightarrow & 236 & \rightarrow & 223 & \rightarrow & 202 \rightarrow 169 \rightarrow 114 \end{array} $
	ω_2	
	ω_3	
	ω_4	
	ω_5	
	ω_6	
	ω_7	$ \begin{array}{ccccccccccc} 248 & \rightarrow & 247 & \rightarrow & 244 & \rightarrow & 239 & \rightarrow & 232 & & \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\ 247 & \rightarrow & 246 & \rightarrow & 243 & \rightarrow & 238 & \rightarrow & 231 & \rightarrow & 220 \rightarrow 199 \rightarrow 156 \end{array} $

Table 1: (continued)

Group	Module	Orbits
	ω_8	

Table 2: Normality and smoothness

Group	Module	L	L -weights	Normality	Smoothness
$A_l = \mathrm{SL}_n$ $n = l + 1$	ω_1, ω_l	GL_l	π'_1	Yes	Yes
	ω_k $1 < k < l$	$\mathrm{PG}(\mathrm{End}_k \times \mathrm{End}_{n-k})$	$\pi'_i + \pi''_i$ $i \leq k, n - k$	Yes	No
	Ad	$\mathbb{C}^\times \times \mathrm{GL}_{n-2}$	$\varepsilon', 2\varepsilon', \pi''_1, \varepsilon' + \pi''_1,$ $\varepsilon' + \pi''_{n-3} - \pi''_{n-2},$ $2\varepsilon' + \pi''_{n-3} - \pi''_{n-2},$ $\varepsilon' + \pi''_1 + \pi''_{n-3} - \pi''_{n-2}$	Yes	$n \leq 3$
$B_l = \mathrm{SO}_n$ $n = 2l + 1$	ω_k $k < l$	$\mathrm{GL}_k \times \mathrm{SO}_{n-2k}$	$\pi'_i + \pi'_j + \pi''_m$ $m \leq i - j$ $m \equiv i - j \pmod{2}$	No $\nexists \pi'_1, \exists 2\pi'_1$	No
	ω_l	GL_l	$\pi'_1, \dots, \pi'_l$	Yes	Yes
$C_l = \mathrm{Sp}_n$ $n = 2l$	ω_k $k < l$	$\mathrm{P}(\mathrm{GL}_k \times \mathrm{Sp}_{n-2k})$	$\pi'_1 + \pi''_1, \dots$	Yes	$n = 4$
	ω_l	$\mathrm{GL}_l/\mathbb{Z}_2$	$2\pi'_1, \dots, 2\pi'_l$	$n = 2$ $\nexists \pi'_2, \exists 2\pi'_2$	$n = 2$
	Ad	$\mathrm{P}(\mathbb{C}^\times \times \mathrm{Sp}_{n-2})$	$\varepsilon' + \pi''_1, 2\varepsilon', 4\varepsilon',$ $2\varepsilon' + 2\pi''_1, 3\varepsilon' + \pi''_1$	Yes	$n \leq 4$
$D_l = \mathrm{SO}_n$ $n = 2l$	ω_k $k \leq l - 2$	$\mathrm{P}(\mathrm{GL}_k \times \mathrm{SO}_{n-2k})$	Same as for B_l , and obtained by $\varepsilon''_l \leftrightarrow \varepsilon''_{l+1}$	Yes	No
	ω_{l-1}, ω_l	$\mathrm{GL}_l/\mathbb{Z}_2$	$\pi'_2, \pi'_4, \dots$	Yes	$n = 6$
G_2	ω_1	GL_2	$\pi'_1, \pi'_2, \pi'_1 + \pi'_2, 2\pi'_2$	Yes	Yes
	ω_2	GL_2	$2\pi'_1, 3\pi'_1, \pi'_2, 2\pi'_2$ $3\pi'_1 - \pi'_2$	No $\nexists \pi'_1, \exists 2\pi'_1$	No
F_4	ω_1	$\mathbb{C}^\times \times \mathrm{Spin}_7$	$\varepsilon' + \omega'_3, 2\varepsilon', 4\varepsilon',$ $2\varepsilon' + \omega'_1, 3\varepsilon' + \omega'_3$	Yes	No
	ω_2	$\mathrm{GL}_2 \times \mathrm{SL}_3$	$\pi'_1 + \pi''_1, \dots$	Yes	No
	ω_3	$\mathrm{SL}_3 \times \mathrm{GL}_2$	$2\pi'_1 + \pi''_1, \dots$	No $\nexists \pi'_2 + \pi''_1$ $\exists 2\pi'_2 + 2\pi''_1$	No
	ω_4	$\mathbb{C}^\times \times \mathrm{Sp}_6$	$\varepsilon' + \omega'_3, 2\varepsilon', 4\varepsilon',$ $2\varepsilon' + 2\omega''_1, 3\varepsilon' + \omega''_3$	No $\nexists \varepsilon' + \omega''_1$ $\exists 2\varepsilon' + 2\omega''_1$	No
E_6	ω_1, ω_5	$\mathbb{C}^\times \times \mathrm{Spin}_{10}$	$\varepsilon' + \omega''_5, 2\varepsilon' + \omega''_1$	Yes	No
	ω_2, ω_4	$\mathrm{GL}_2 \times \mathrm{SL}_5$	$\pi'_1 + \pi''_2, \dots$	Yes	No
	ω_3	$\mathrm{GL}_2 \times \mathrm{SL}_3 \times \mathrm{SL}_3$	$\pi'_1 + \pi''_1 + \pi'''_1, \dots$	Yes	No

Table 2: (continued)

Group	Module	L	L -weights	Normality	Smoothness
	ω_6	$\mathrm{GL}_6/\mathbb{Z}_3$	$\pi'_3, \pi'_6, 2\pi'_6,$ $\pi'_3 + \pi'_6, \pi'_1 + \pi'_5$	Yes	No
E_7	ω_1	$\mathbb{C}^\times \times E_6$	$\varepsilon' + \omega''_1, 2\varepsilon' + \omega''_5, 3\varepsilon'$	Yes	No
	ω_2	$\mathrm{GL}_2 \times \mathrm{Spin}_{10}$	$\pi'_1 + \omega''_5, \dots$	Yes	No
	ω_3	$\mathrm{GL}_3 \times \mathrm{SL}_5$	$\pi'_1 + \pi''_2, \dots$	Yes	No
	ω_4	$\mathrm{GL}_2 \times \mathrm{SL}_3 \times \mathrm{SL}_4$	$\pi'_1 + \pi''_1 + \pi'''_1, \dots$	Yes	No
	ω_5	$\mathrm{GL}_2 \times \mathrm{SL}_6$	$\pi'_1 + \pi''_2, \dots$	Yes	No
	ω_6	$\mathbb{C}^\times \times \mathrm{Spin}_{12}/\mathbb{Z}_2$	$\varepsilon' + \omega''_6, 2\varepsilon', 4\varepsilon',$ $2\varepsilon' + \omega''_2, 3\varepsilon' + \omega''_6$	Yes	No
	ω_7	$\mathrm{GL}_7/\mathbb{Z}_3$	$\pi'_3, \dots$	Yes	No
E_8	ω_1	$\mathbb{C}^\times \times E_7$	$\varepsilon' + \omega''_1, 2\varepsilon', 4\varepsilon',$ $2\varepsilon' + \omega''_6, 3\varepsilon' + \omega''_1$	Yes	No
	ω_2	$\mathrm{GL}_2 \times E_6$	$\pi'_1 + \omega''_1, \dots$	Yes	No
	ω_3	$\mathrm{GL}_3 \times \mathrm{Spin}_{10}$	$\pi'_1 + \omega''_5, \dots$	Yes	No
	ω_4	$\mathrm{GL}_4 \times \mathrm{SL}_5$	$\pi'_1 + \pi''_2, \dots$	Yes	No
	ω_5	$\mathrm{GL}_2 \times \mathrm{SL}_3 \times \mathrm{SL}_5$	$\pi'_1 + \pi''_1 + \pi'''_1, \dots$	Yes	No
	ω_6	$\mathrm{GL}_2 \times \mathrm{SL}_7$	$\pi'_1 + \pi''_2, \dots$	Yes	No
	ω_7	$\mathbb{C}^\times \times \mathrm{Spin}_{14}$	$\varepsilon' + \omega''_7, \dots$	Yes	No
	ω_8	$\mathrm{GL}_8/\mathbb{Z}_3$	$\pi'_3, \dots$	Yes	No

REFERENCES

- [AB1] V. Alexeev, M. Brion, *Stable reductive varieties I: Affine varieties*, preprint, 2002, [arXiv:math.AG/0207272](http://xxx.lanl.gov), <http://xxx.lanl.gov>.
- [AB2] V. Alexeev, M. Brion, *Stable reductive varieties II: Projective case*, preprint, 2002, [arXiv:math.AG/0207274](http://xxx.lanl.gov), <http://xxx.lanl.gov>.
- [BCP] E. Bifet, C. de Concini, C. Procesi, *Cohomology of regular embeddings*, Adv. Math. **82** (1990), no. 1, 1–34.
- [Br1] M. Brion, *Groupe de Picard et nombres caractéristiques des variétés sphériques*, Duke Math. J. **58** (1989), no. 2, 397–424.
- [Br2] M. Brion, *Sur la géométrie des variétés sphériques*, Comment. Math. Helv. **66** (1991), 237–262.
- [Br3] M. Brion, *Variétés sphériques*, Notes de la session de la S. M. F. “Opérations hamiltoniennes et opérations de groupes algébriques”, Grenoble, 1997. <http://www-fourier.ujf-grenoble.fr/~mbrion/spheriques.ps>
- [Br4] M. Brion, *The behaviour at infinity of the Bruhat decomposition*, Comment. Math. Helv. **73** (1998), 137–174.
- [BI] M. Brion, S. P. Inamdar, *Frobenius splitting of spherical varieties*, Algebraic groups and their generalizations: Classical methods (W. J. Haboush, B. J. Parshall, eds.), Proc. Symp. Pure Math., vol. 56, part 1, pp. 207–218, AMS, Providence, 1994.
- [BLV] M. Brion, D. Luna, Th. Vust, *Espaces homogènes sphériques*, Invent. Math. **84** (1986), 617–632.
- [BP] M. Brion, P. Polo, *Large Schubert varieties*, Representation Theory **4** (2000), 97–126.
- [Con] C. de Concini, *Normality and non-normality of certain semigroups and orbit closures*, preprint, 2003.
- [CP-I] C. de Concini, C. Procesi, *Complete symmetric varieties*, Lect. Notes in Math., vol. 996, 1983, pp. 1–44.

- [CP-II] C. de Concini, C. Procesi, *Complete symmetric varieties, II*, Algebraic groups and related topics (R. Hotta, ed.), Adv. Studies in Pure Math., no. 6, pp. 481–513, Kinokuniya, Tokyo, 1985.
- [CP] C. de Concini, C. Procesi, *Cohomology of compactifications of algebraic groups*, Duke Math. J. **53** (1986), 585–594.
- [Fal] G. Faltings, *Explicit resolution of local singularities of moduli-spaces*, J. Reine Angew. Math. **483** (1997), 183–196.
- [Kan] S. S. Kannan, *Projective normality of the wonderful compactification of semisimple adjoint groups*, Math. Z. **239** (2002), 673–682.
- [Kap] M. M. Kapranov, *Hypergeometric functions on reductive groups*, Integrable systems and algebraic geometry (M.-H. Saito, ed.), pp. 236–281, World Scientific, Singapore, 1998.
- [Kn] F. Knop, *The Luna–Vust theory of spherical embeddings*, Proc. Hyderabad Conf. on Algebraic Groups (S. Ramanan, ed.), pp. 225–249, Manoj Prakashan, Madras, 1991.
- [Kr] H. Kraft, *Geometrische Methoden in der Invariantentheorie*, Vieweg, Braunschweig–Wiesbaden, 1985.
- [LP] P. Littelmann, C. Procesi, *Equivariant cohomology of wonderful compactifications*, Operator algebras, unitary representations, enveloping algebras, and Invariant Theory, (Paris, 1989), Progress in Math., vol. 92, pp. 219–262, Birkhäuser, Boston, 1990.
- [Lu] D. Luna, *Slices étales*, Mém. Soc. Math. France (N.S.) **33** (1973), 81–105.
- [LV] D. Luna, Th. Vust, *Plongements d’espaces homogènes*, Comment. Math. Helv. **58** (1983), 186–245.
- [Ner] Yu. A. Neretin, *Hinges and the Study–Semple–Satake–Furstenberg–De Concini–Procesi–Oshima boundary*, Kirillov’s Seminar on Representation Theory, AMS Transl. (2) **181** (1998), 165–230.
- [Po] V. L. Popov, *Contractions of the actions of reductive algebraic groups*, Mat. Sb. (N.S.) **130** (1986), no. 3, 310–334.
English translation: Math. USSR-Sb. **58** (1987), no. 2, 311–335.
- [Pu1] M. S. Putcha, *Linear algebraic monoids*, Cambridge University Press, 1988.
- [Pu2] M. S. Putcha, *Monoids on groups with BN pair*, J. Algebra **120** (1989), 139–169.
- [PR] M. S. Putcha, L. E. Renner, *The system of idempotents and the lattice of $\mathcal{J}$ -classes of reductive algebraic monoids*, J. Algebra **116** (1988), 385–399.
- [Re1] L. E. Renner, *Classification of semisimple algebraic monoids*, Trans. Amer. Math. Soc. **292** (1985), 193–223.
- [Re2] L. E. Renner, *Classification of semisimple varieties*, J. Algebra **122** (1989), 275–287.
- [Rit] A. Rittatore, *Algebraic monoids and group embeddings*, Transformation Groups **3** (1998), 375–396.
- [Sem] J. G. Semple, *The variety whose points represent complete collineations of S_r on S'_r* , Rend. Mat. Univ. Roma (5), **10** (1951), 201–208.
- [St1] E. S. Strickland, *A vanishing theorem for group compactifications*, Math. Ann. **277** (1987), 165–171.
- [St2] E. S. Strickland, *Computing the equivariant cohomology of group compactifications*, Math. Ann. **291** (1991), no. 2, 275–280.
- [Vin] E. B. Vinberg, *On reductive algebraic semigroups*, Lie Groups and Lie Algebras: E. B. Dynkin Seminar (S. Gindikin, E. Vinberg, eds.), AMS Transl. (2) **169** (1995), 145–182.
- [VO] E. B. Vinberg, A. L. Onishchik, *Seminar on Lie groups and algebraic groups*, Nauka, Moscow, 1988.

English translation: *Lie groups and algebraic groups*, Springer, Berlin–Heidelberg–New York, 1990.

- [Vu] T. Vust, *Plongements d'espaces symétriques algébriques: une classification*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **XVII** (1990), no. 2, 165–194.

DEPARTMENT OF HIGHER ALGEBRA, FACULTY OF MECHANICS AND MATHEMATICS, MOSCOW STATE UNIVERSITY, 119992 MOSCOW, RUSSIA

E-mail address: `timashev@mech.math.msu.su`